

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number  
**WO 01/94585 A1**(51) International Patent Classification<sup>7</sup>: **C12N 15/13**,  
C07K 16/24, 16/46, A61K 47/48, C07K 19/00, C12N  
15/62, 15/70, 1/21, A61K 39/395, A61P 19/02, 37/06John [GB/GB]; 208 Bath Road, Slough, Berkshire SL1  
3WE (GB).

(21) International Application Number: PCT/GB01/02477

(74) Agents: **MERCER, Christopher, Paul** et al.; Carpmals  
& Ransford, 43 Bloomsbury Square, London WC1A 2RA  
(GB).

(22) International Filing Date: 5 June 2001 (05.06.2001)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:  
0013810.7 6 June 2000 (06.06.2000) GB(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,  
ZW.(71) Applicant (*for all designated States except US*): **CELL-  
TECH R & D LIMITED** [GB/GB]; 208 Bath Road,  
Slough, Berkshire SL1 3WE (GB).(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ATHWAL, Diljeet**,  
Singh [GB/GB]; 208 Bath Road, Slough, Berkshire SL1  
3WE (GB). **BROWN, Derek, Thomas** [GB/GB]; 208 Bath  
Road, Slough, Berkshire SL1 3WE (GB). **WEIR, Andrew**,  
Neil, Charles [GB/GB]; 208 Bath Road, Slough, Berks-  
hire SL1 3WE (GB). **POPPELWELL, Andrew, George**  
[GB/GB]; 208 Bath Road, Slough, Berkshire SL1 3WE  
(GB). **CHAPMAN, Andrew, Paul** [GB/GB]; 208 Bath  
Road, Slough, Berkshire SL1 3WE (GB). **KING, David**,

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.(54) Title: ANTIBODY MOLECULES HAVING SPECIFICITY FOR HUMAN TUMOR NECROSIS FACTOR ALPHA, AND  
USE THEREOF(57) Abstract: There is disclosed antibody molecules containing at least one CDR derived from a mouse monoclonal antibody having specificity for human TNF $\alpha$ . There is also disclosed a CDR grafted antibody wherein at least one of the CDRs is a hybrid CDR. Further disclosed are DNA sequences encoding the chains of the antibody molecules, vectors, transformed host cells and uses of the antibody molecules in the treatment of diseases mediated by TNF $\alpha$ .

## ANTIBODY MOLECULES HAVING SPECIFICITY FOR HUMAN TUMOR NECROSIS FACTOR ALPHA, AND USE THEREOF

The present invention relates to an antibody molecule having specificity for antigenic determinants of human tumour necrosis factor alpha (TNF $\alpha$ ). The present invention also relates to the therapeutic uses of the antibody molecule and methods for producing the antibody molecule.

This invention relates to antibody molecules. In an antibody molecule, there are two heavy chains and two light chains. Each heavy chain and each light chain has at its N-terminal end a variable domain. Each variable domain is composed of four framework regions (FRs) alternating with three complementarily determining regions (CDRs). The residues in the variable domains are conventionally numbered according to a system devised by Kabat *et al.* This system is set forth in Kabat *et al.*, 1987, in Sequences of Proteins of Immunological Interest, US Department of Health and Human Services, NIH, USA (hereafter "Kabat *et al. (supra)*"). This numbering system is used in the present specification except where otherwise indicated.

The Kabat residue designations do not always correspond directly with the linear numbering of the amino acid residues. The actual linear amino acid sequence may contain fewer or additional amino acids than in the strict Kabat numbering corresponding to a shortening of, or insertion into, a structural component, whether framework or CDR, of the basic variable domain structure. The correct Kabat numbering of residues may be determined for a given antibody by alignment of residues of homology in the sequence of the antibody with a "standard" Kabat numbered sequence.

The CDRs of the heavy chain variable domain are located at residues 31-35 (CDRH1), residues 50-65 (CDRH2) and residues 95-102 (CDRH3) according to the Kabat numbering.

The CDRs of the light chain variable domain are located at residues 24-34 (CDRL1), residues 50-56 (CDRL2) and residues 89-97 (CDRL3) according to the Kabat numbering.

Construction of CDR-grafted antibodies is described in European Patent Application EP-A-0239400, which discloses a process in which the CDRs of a mouse monoclonal antibody are grafted onto the framework regions of the variable domains of a human immunoglobulin by site directed mutagenesis using long oligonucleotides. The

CDRs determine the antigen binding specificity of antibodies and are relatively short peptide sequences carried on the framework regions of the variable domains.

The earliest work on humanising monoclonal antibodies by CDR-grafting was carried out on monoclonal antibodies recognising synthetic antigens, such as NP.  
5 However, examples in which a mouse monoclonal antibody recognising lysozyme and a rat monoclonal antibody recognising an antigen on human T-cells were humanised by CDR-grafting have been described by Verhoeyen *et al.* (Science, 239, 1534-1536, 1988) and Riechmann *et al.* (Nature, 332, 323-324, 1988), respectively.

Riechmann *et al.*, found that the transfer of the CDRs alone (as defined by Kabat  
10 (Kabat *et al.* (*supra*) and Wu *et al.*, J. Exp. Med., 132, 211-250, 1970)) was not sufficient to provide satisfactory antigen binding activity in the CDR-grafted product. It was found that a number of framework residues have to be altered so that they correspond to those of the donor framework region. Proposed criteria for selecting which framework residues need to be altered are described in International Patent Application WO 90/07861.

15 A number of reviews discussing CDR-grafted antibodies have been published, including Vaughan *et al.* (Nature Biotechnology, 16, 535-539, 1998).

TNF $\alpha$  is a pro-inflammatory cytokine that is released by and interacts with cells of the immune system. Thus, TNF $\alpha$  is released by macrophages that have been activated by lipopolysaccharides (LPS) of gram negative bacteria. As such, TNF $\alpha$  appears to be an  
20 endogenous mediator of central importance involved in the development and pathogenesis of endotoxic shock associated with bacterial sepsis. TNF $\alpha$  has also been shown to be up-regulated in a number of human diseases, including chronic diseases such as rheumatoid arthritis, Crohn's disease, ulcerative colitis and multiple sclerosis. Mice transgenic for human TNF $\alpha$  produce high levels of TNF $\alpha$  constitutively and develop a spontaneous,  
25 destructive polyarthritis resembling rheumatoid arthritis (Kaffer *et al.*, EMBO J., 10, 4025-4031, 1991). TNF $\alpha$  is therefore referred to as a pro-inflammatory cytokine.

Monoclonal antibodies against TNF $\alpha$  have been described in the prior art. Meager *et al.*, (Hybridoma, 6, 305-311, 1987) describe murine monoclonal antibodies against recombinant TNF $\alpha$ . Fendly *et al.*, (Hybridoma, 6, 359-370, 1987) describe the use of  
30 murine monoclonal antibodies against recombinant TNF $\alpha$  in defining neutralising epitopes on TNF. Shimamoto *et al.*, (Immunology Letters, 17, 311-318, 1988) describe the use of murine monoclonal antibodies against TNF $\gamma$  and their use in preventing endotoxic shock in mice. Furthermore, in International Patent Application WO 92/11383, recombinant

antibodies, including CDR-grafted antibodies, specific for TNF $\alpha$  are disclosed. Rankin *et al.*, (British J. Rheumatology, 34, 334-342, 1995) describe the use of such CDR-grafted antibodies in the treatment of rheumatoid arthritis. US-A-5 919 452 discloses anti-TNF chimeric antibodies and their use in treating pathologies associated with the presence of

5 TNF.

Antibodies to TNF $\alpha$  have been proposed for the prophylaxis and treatment of endotoxic shock (Beutler *et al.*, Science, 234, 470-474, 1985). Bodmer *et al.*, (Critical Care Medicine, 21, S441-S446, 1993) and Wherry *et al.*, (Critical Care Medicine, 21, S436-S440, 1993) discuss the therapeutic potential of anti-TNF $\alpha$  antibodies in the treatment of

10 septic shock. The use of anti-TNF $\alpha$  antibodies in the treatment of septic shock is also discussed by Kirschenbaum *et al.*, (Critical Care Medicine, 26, 1625-1626, 1998). Collagen-induced arthritis can be treated effectively using an anti-TNF $\alpha$  monoclonal antibody (Williams *et al.* (PNAS-USA, 89, 9784-9788, 1992)).

Increased levels of TNF $\alpha$  are found in both the synovial fluid and peripheral blood

15 of patients suffering from rheumatoid arthritis. When TNF $\alpha$  blocking agents are administered to patients suffering from rheumatoid arthritis, they reduce inflammation, improve symptoms and retard joint damage (McKown *et al.* (Arthritis Rheum., 42, 1204-1208, 1999).

The use of anti-TNF $\alpha$  antibodies in the treatment of rheumatoid arthritis and

20 Crohn's disease is discussed in Feldman *et al.*, (Transplantation Proceedings, 30, 4126-4127, 1998), Adorini *et al.*, (Trends in Immunology Today, 18, 209-211, 1997) and in Feldman *et al.*, (Advances in Immunology, 64, 283-350, 1997). The antibodies to TNF $\alpha$  used in such treatments are generally chimeric antibodies, such as those described in US-A-5 919 452.

25 Two TNF $\alpha$  blocking products are currently licensed for the treatment of rheumatoid arthritis. The first, called etanercept, is marketed by Immunex Corporation as Enbrel™. It is a recombinant fusion protein comprising two p75 soluble TNF-receptor domains linked to the Fc portion of a human immunoglobulin. The second, called infliximab, is marketed by Centocor Corporation as Remicade™. It is a chimeric antibody having murine anti-

30 TNF $\alpha$  variable domains and human IgG1 constant domains.

The prior art recombinant anti-TNF $\alpha$  antibody molecules generally have a reduced affinity for TNF $\alpha$  compared to the antibodies from which the variable regions or CDRs are

derived, generally have to be produced in mammalian cells and are expensive to manufacture. Prior art anti-TNF $\alpha$  antibodies are described in Stephens *et al.*, (Immunology, 85; 668-674, 1995), GB-A-2 246 570 and GB-A-2 297 145.

There is a need for an antibody molecule to treat chronic inflammatory diseases  
5 which can be used repeatedly and produced easily and efficiently. There is also a need for an antibody molecule which has high affinity for TNF $\alpha$  and low immunogenicity in humans.

In a first aspect, the present invention provides an antibody molecule having specificity for TNF $\alpha$ , comprising a heavy chain wherein the variable domain comprises a  
10 CDR (as defined by Kabat *et al.*, (*supra*)) having the sequence given as H1 in Figure 3 (SEQ ID NO:1) for CDRH1, as H2' in Figure 3 (SEQ ID NO:2) or as H2 in Figure 3 (SEQ ID NO:7) for CDRH2 or as H3 in Figure 3 (SEQ ID NO:3) for CDRH3.

The antibody molecule of the first aspect of the present invention comprises at least one CDR selected from H1, H2' or H2 and H3 (SEQ ID NO:1; SEQ ID NO:2 or SEQ ID  
15 NO:7 and SEQ ID NO:3) for the heavy chain variable domain. Preferably, the antibody molecule comprises at least two and more preferably all three CDRs in the heavy chain variable domain.

In a second aspect of the present invention, there is provided an antibody molecule having specificity for human TNF $\alpha$ , comprising a light chain wherein the variable domain  
20 comprises a CDR (as defined by Kabat *et al.*, (*supra*)) having the sequence given as L1 in Figure 3 (SEQ ID NO:4) for CDRL1, L2 in Figure 3 (SEQ ID NO:5) for CDRL2 or L3 in Figure 3 (SEQ ID NO:6) for CDRL3.

The antibody molecule of the second aspect of the present invention comprises at least one CDR selected from L1, L2 and L3 (SEQ ID NO:4 to SEQ ID NO:6) for the light  
25 chain variable domain. Preferably, the antibody molecule comprises at least two and more preferably all three CDRs in the light chain variable domain.

The antibody molecules of the first and second aspects of the present invention preferably have a complementary light chain or a complementary heavy chain, respectively.

Preferably, the antibody molecule of the first or second aspect of the present  
30 invention comprises a heavy chain wherein the variable domain comprises a CDR (as defined by Kabat *et al.*, (*supra*)) having the sequence given as H1 in Figure 3 (SEQ ID NO:1) for CDRH1, as H2' or H2 in Figure 3 (SEQ ID NO:2 or SEQ ID NO:7) for CDRH2 or as H3 in Figure 3 (SEQ ID NO:3) for CDRH3 and a light chain wherein the variable

domain comprises a CDR (as defined by Kabat *et al.*, (*supra*)) having the sequence given as L1 in Figure 3 (SEQ ID NO:4) for CDRL1, as L2 in Figure 3 (SEQ ID NO:5) for CDRL2 or as L3 in Figure 3 (SEQ ID NO:6) for CDRL3.

The CDRs given in SEQ IDS NOS:1 and 3 to 7 and in Figure 3 referred to above  
5 are derived from a mouse monoclonal antibody hTNF40. However, SEQ ID NO:2 consists of a hybrid CDR. The hybrid CDR comprises part of heavy chain CDR2 from mouse monoclonal antibody hTNF40 (SEQ ID NO:7) and part of heavy chain CDR2 from a human group 3 germline V region sequence.

The complete sequences of the variable domains of the mouse hTNF40 antibody  
10 are shown in Figures 6 (light chain) (SEQ ID NO:99) and Figure 7 (heavy chain) (SEQ ID NO:100). This mouse antibody is referred to below as "the donor antibody".

A first alternatively preferred embodiment of the first or second aspect of the present invention is the mouse monoclonal antibody hTNF40 having the light and heavy chain variable domain sequences shown in Figure 6 (SEQ ID NO:99) and Figure 7 (SEQ  
15 ID NO:100), respectively. The light chain constant region of hTNF40 is kappa and the heavy chain constant region is IgG2a.

In a second alternatively preferred embodiment, the antibody according to either of the first and second aspects of the present invention is a chimeric mouse/human antibody molecule, referred to herein as the chimeric hTNF40 antibody molecule. The chimeric  
20 antibody molecule comprises the variable domains of the mouse monoclonal antibody hTNF40 (SEQ ID NOS:99 and 100) and human constant domains. Preferably, the chimeric hTNF40 antibody molecule comprises the human C kappa domain (Hieter *et al.*, Cell, 22, 197-207, 1980; Genebank accession number J00241) in the light chain and the human gamma 4 domains (Flanagan *et al.*, Nature, 300, 709-713, 1982) in the heavy chain.

25 In a third alternatively preferred embodiment, the antibody according to either of the first and second aspects of the present invention is a CDR-grafted antibody molecule. The term "a CDR-grafted antibody molecule" as used herein refers to an antibody molecule wherein the heavy and/or light chain contains one or more CDRs (including, if desired, a hybrid CDR) from the donor antibody (e.g. a murine monoclonal antibody) grafted into a  
30 heavy and/or light chain variable region framework of an acceptor antibody (e.g. a human antibody).

Preferably, such a CDR-grafted antibody has a variable domain comprising human acceptor framework regions as well as one or more of the donor CDRs referred to above.

When the CDRs are grafted, any appropriate acceptor variable region framework sequence may be used having regard to the class/type of the donor antibody from which the CDRs are derived, including mouse, primate and human framework regions. Examples of human frameworks which can be used in the present invention are KOL, NEWM, REI, EU, 5 TUR, TEI, LAY and POM (Kabat *et al. (supra)*). For example, KOL and NEWM can be used for the heavy chain, REI can be used for the light chain and EU, LAY and POM can be used for both the heavy chain and the light chain. The preferred framework regions for the light chain are the human group 1 framework regions shown in Figure 1 (SEQ ID NOS:83, 85, 87 and 89). The preferred framework regions for the heavy chain are the 10 human group 1 and group 3 framework regions shown in Figure 2 (SEQ ID NOS:91, 93, 95 and 97 and SEQ ID NOS:106, 107, 108 and 109), respectively.

In a CDR-grafted antibody of the present invention, it is preferred to use as the acceptor antibody one having chains which are homologous to the chains of the donor antibody. The acceptor heavy and light chains do not necessarily need to be derived from 15 the same antibody and may, if desired, comprise composite chains having framework regions derived from different chains.

Also, in a CDR-grafted antibody of the present invention, the framework regions need not have exactly the same sequence as those of the acceptor antibody. For instance, unusual residues may be changed to more frequently-occurring residues for that acceptor 20 chain class or type. Alternatively, selected residues in the acceptor framework regions may be changed so that they correspond to the residue found at the same position in the donor antibody. Such changes should be kept to the minimum necessary to recover the affinity of the donor antibody. A protocol for selecting residues in the acceptor framework regions which may need to be changed is set forth in WO 91/09967.

25 Preferably, in a CDR-grafted antibody molecule of the present invention, if the acceptor heavy chain has human group 1 framework regions (shown in Figure 2) (SEQ ID NOS:91, 93, 95 and 97), then the acceptor framework regions of the heavy chain comprise, in addition to one or more donor CDRs, donor residues at positions 28, 69 and 71 (according to Kabat *et al. (supra)*).

30 Alternatively, if the acceptor heavy chain has group 1 framework regions, then the acceptor framework regions of the heavy chain comprise, in addition to one or more donor CDRs, donor residues at positions 28, 38, 46, 67, 69 and 71 (according to Kabat *et al. (supra)*).

Preferably, in a CDR-grafted antibody molecule of the present invention, if the acceptor heavy chain has human group 3 framework regions (shown in Figure 2) (SEQ ID NOS:106, 107, 108 and 109), then the acceptor framework regions of the heavy chain comprise, in addition to one or more donor CDRs, donor residues at positions 27, 28, 30, 48, 49, 69, 71, 73, 76 and 78 (according to Kabat *et al. (supra)*).

Preferably, in a CDR-grafted antibody molecule according to the present invention, if the acceptor light chain has human group 1 framework regions (shown in Figure 1) (SEQ ID NOS:83, 85, 87 and 89) then the acceptor framework regions of the light chain comprise donor residues at positions 46 and 60 (according to Kabat *et al. (supra)*).

Donor residues are residues from the donor antibody, i.e. the antibody from which the CDRs were originally derived.

The antibody molecule of the present invention may comprise: a complete antibody molecule, having full length heavy and light chains; a fragment thereof, such as a Fab, modified Fab, Fab', F(ab')<sub>2</sub> or Fv fragment; a light chain or heavy chain monomer or dimer; a single chain antibody, e.g. a single chain Fv in which the heavy and light chain variable domains are joined by a peptide linker. Similarly, the heavy and light chain variable regions may be combined with other antibody domains as appropriate.

Preferably the antibody molecule of the present invention is a Fab fragment. Preferably the Fab fragment has a heavy chain having the sequence given as SEQ ID NO:111 and a light chain having the sequence given as SEQ ID NO:113. The amino acid sequences given in SEQ ID NO:111 and SEQ ID NO:113 are preferably encoded by the nucleotide sequences given in SEQ ID NO:110 and SEQ ID NO:112, respectively.

Alternatively, it is preferred that the antibody molecule of the present invention is a modified Fab fragment wherein the modification is the addition to the C-terminal end of its heavy chain one or more amino acids to allow the attachment of an effector or reporter molecule. Preferably, the additional amino acids form a modified hinge region containing one or two cysteine residue to which the effector or reporter molecule may be attached. Such a modified Fab fragment preferably has a heavy chain having the sequence given as SEQ ID NO:115 and the light chain having the sequence given as SEQ ID NO:113. The amino acid sequence given in SEQ ID NO:115 is preferably encoded by the nucleotide sequence given in SEQ ID NO:114.

A preferred effector group is a polymer molecule, which may be attached to the modified Fab fragment to increase its half-life *in vivo*.



The polymer molecule may, in general, be a synthetic or a naturally occurring polymer, for example an optionally substituted straight or branched chain polyalkylene, polyalkenylene or polyoxyalkylene polymer or a branched or unbranched polysaccharide, e.g. a homo- or hetero- polysaccharide.

5 Particular optional substituents which may be present on the above-mentioned synthetic polymers include one or more hydroxy, methyl or methoxy groups. Particular examples of synthetic polymers include optionally substituted straight or branched chain poly(ethyleneglycol), poly(propyleneglycol) poly(vinylalcohol) or derivatives thereof, especially optionally substituted poly(ethyleneglycol) such as methoxypoly(ethyleneglycol)  
10 or derivatives thereof. Particular naturally occurring polymers include lactose, amylose, dextran, glycogen or derivatives thereof. "Derivatives" as used herein is intended to include reactive derivatives, for example thiol-selective reactive groups such as maleimides and the like. The reactive group may be linked directly or through a linker segment to the polymer. It will be appreciated that the residue of such a group will in some instances form  
15 part of the product as the linking group between the antibody fragment and the polymer.

The size of the polymer may be varied as desired, but will generally be in an average molecular weight range from 500Da to 50000Da, preferably from 5000 to 4000Da and more preferably from 25000 to 4000Da. The polymer size may in particular be selected on the basis of the intended use of the product. Thus, for example, where the  
20 product is intended to leave the circulation and penetrate tissue, for example for use in the treatment of a tumour, it may be advantageous to use a small molecular weight polymer, for example with a molecular weight of around 5000Da. For applications where the product remains in the circulation, it may be advantageous to use a higher molecular weight polymer, for example having a molecular weight in the range from 25000Da to 40000Da.

25 Particularly preferred polymers include a polyalkylene polymer, such as a poly(ethyleneglycol) or, especially, a methoxypoly(ethyleneglycol) or a derivative thereof, and especially with a molecular weight in the range from about 25000Da to about 40000Da.

Each polymer molecule attached to the modified antibody fragment may be  
30 covalently linked to the sulphur atom of a cysteine residue located in the fragment. The covalent linkage will generally be a disulphide bond or, in particular, a sulphur-carbon bond.

Where desired, the antibody fragment may have one or more effector or reporter molecules attached to it. The effector or reporter molecules may be attached to the antibody fragment through any available amino acid side-chain or terminal amino acid functional group located in the fragment, for example any free amino, imino, hydroxyl or  
5 carboxyl group.

An activated polymer may be used as the starting material in the preparation of polymer-modified antibody fragments as described above. The activated polymer may be any polymer containing a thiol reactive group such as an  $\alpha$ -halocarboxylic acid or ester, e.g. iodoacetamide, an imide, e.g. maleimide, a vinyl sulphone or a disulphide. Such  
10 starting materials may be obtained commercially (for example from Shearwater Polymers Inc., Huntsville, AL, USA) or may be prepared from commercially available starting materials using conventional chemical procedures.

As regards attaching poly(ethyleneglycol) (PEG) moieties, reference is made to "Poly(ethyleneglycol) Chemistry, Biotechnical and Biomedical Applications", 1992, J.  
15 Milton Harris (ed), Plenum Press, New York, "Poly(ethyleneglycol) Chemistry and Biological Applications", 1997, J. Milton Harris and S. Zalipsky (eds), American Chemical Society, Washington DC and "Bioconjugation Protein Coupling Techniques for the Biomedical Sciences", 1998, M. Aslam and A. Dent, Grove Publishers, New York.

Where it is desired to obtain an antibody fragment linked to an effector or reporter  
20 molecule, this may be prepared by standard chemical or recombinant DNA procedures in which the antibody fragment is linked either directly or via a coupling agent to the effector or reporter molecule either before or after reaction with the activated polymer as appropriate. Particular chemical procedures include, for example, those described in WO 93/62331, WO 92/22583, WO 90,195 and WO 89/1476. Alternatively, where the effector  
25 or reporter molecule is a protein or polypeptide the linkage may be achieved using recombinant DNA procedures, for example as described in WO 86/01533 and EP-A-0392745.

Preferably, the modified Fab fragment of the present invention is PEGylated (i.e. has PEG (poly(ethyleneglycol)) covalently attached thereto) according to the method  
30 disclosed in EP-A-0948544. Preferably the antibody molecule of the present invention is a PEGylated modified Fab fragment as shown in Figure 13. As shown in Figure 13, the modified Fab fragment has a maleimide group covalently linked to a single thiol group in a modified hinge region. A lysine residue is covalently linked to the maleimide group. To

each of the amine groups on the lysine residue is attached a methoxypoly(ethyleneglycol) polymer having a molecular weight of approximately 20,000 Da. The total molecular weight of the entire effector molecule is therefore approximately 40,000 Da.

Preferably, in the compound shown in Figure 13, the heavy chain of the antibody  
5 part has the sequence given as SEQ ID NO:115 and the light chain has the sequence given in SEQ ID NO:113. This compound is referred to herein as CDP870.

The constant region domains of the antibody molecule of the present invention, if present, may be selected having regard to the proposed function of the antibody molecule, and in particular the effector functions which may be required. For example, the constant  
10 region domains may be human IgA, IgD, IgE, IgG or IgM domains. In particular, human IgG constant region domains may be used, especially of the IgG1 and IgG3 isotypes when the antibody molecule is intended for therapeutic uses and antibody effector functions are required. Alternatively, IgG2 and IgG4 isotypes may be used when the antibody molecule is intended for therapeutic purposes and antibody effector functions are not required, e.g.  
15 for simply blocking TNF $\alpha$  activity.

Also, the antibody molecule of the present invention may have an effector or a reporter molecule attached to it. For instance, it may have a macrocycle, for chelating a heavy metal atom, or a toxin, such as ricin, attached to it by a covalent bridging structure. Alternatively, procedures of recombinant DNA technology may be used to produce an  
20 antibody molecule in which the Fc fragment (CH2, CH3 and hinge domains), the CH2 and CH3 domains or the CH3 domain of a complete immunoglobulin molecule has (have) been replaced by, or has attached thereto by peptide linkage, a functional non-immunoglobulin protein, such as an enzyme or toxin molecule.

The antibody molecule of the present invention preferably has a binding affinity of  
25 at least  $0.85 \times 10^{-10}$  M, more preferably at least  $0.75 \times 10^{-10}$  M and most preferably at least  $0.5 \times 10^{-10}$  M. (It is worth noting that the preferred humanised antibody molecule of the present invention, as described below, has an affinity of about  $0.5 \times 10^{-10}$  M, which is better than the affinity of the murine monoclonal antibody from which it is derived. The murine antibody has an affinity of about  $0.85 \times 10^{-10}$  M.)

30 Preferably, the antibody molecule of the present invention comprises the light chain variable domain hTNF40-gL1 (SEQ ID NO:8) and the heavy chain variable domain gh3hTNF40.4 (SEQ ID NO:11). The sequences of the variable domains of these light and heavy chains are shown in Figures 8 and 11, respectively.

The present invention also relates to variants of the antibody molecule of the present invention, which have an improved affinity for TNF $\alpha$ . Such variants can be obtained by a number of affinity maturation protocols including mutating the CDRs (Yang *et al.*, J. Mol. Biol., 254, 392-403, 1995), chain shuffling (Marks *et al.*, Bio/Technology, 10, 779-783, 1992), use of mutator strains of *E. coli* (Low *et al.*, J. Mol. Biol., 250, 359-368, 1996), DNA shuffling (Patten *et al.*, Curr. Opin. Biotechnol., 8, 724-733, 1997), phage display (Thompson *et al.*, J. Mol. Biol., 256, 77-88, 1996) and sexual PCR (Cramer *et al.*, Nature, 391, 288-291, 1998). Vaughan *et al.* (*supra*) discusses these methods of affinity maturation.

10 The present invention also provides a DNA sequence encoding the heavy and/or light chain(s) of the antibody molecule of the present invention.

Preferably, the DNA sequence encodes the heavy or the light chain of the antibody molecule of the present invention.

In one preferred embodiment, the DNA sequence encodes a light chain and 15 comprises the sequence shown in SEQ ID NO:8 (hTNF40-gL1) or SEQ ID NO:9 (h-TNF-40-gL2) or a degenerate equivalent thereof.

In an alternatively preferred embodiment, the DNA sequence encodes a heavy chain and comprises the sequence shown in SEQ ID NO:10 (gh1hTNF40.4) or SEQ ID NO:11 (gh3hTNF40.4) or a degenerate equivalent thereof.

20 The DNA sequence of the present invention may comprise synthetic DNA, for instance produced by chemical processing, cDNA, genomic DNA or any combination thereof.

The present invention also relates to a cloning or expression vector comprising one or more DNA sequences of the present invention. Preferably, the cloning or expression 25 vector comprises two DNA sequences, encoding the light chain and the heavy chain of the antibody molecule of the present invention, respectively.

In a preferred embodiment, the present invention provides an *E. coli* expression vector comprising a DNA sequence of the present invention. Preferably the expression vector is pTTO(CDP870) as shown schematically in Figure 22.

30 The present invention also comprises vector pDNAbEng-G1 as shown in Figure 19.

General methods by which the vectors may be constructed, transfection methods and culture methods are well known to those skilled in the art. In this respect, reference is made to "Current Protocols in Molecular Biology", 1999, F. M. Ausubel (ed), Wiley

Interscience, New York and the Maniatis Manual produced by Cold Spring Harbor Publishing.

DNA sequences which encode the antibody molecule of the present invention can be obtained by methods well known to those skilled in the art. For example, DNA  
5 sequences coding for part or all of the antibody heavy and light chains may be synthesised as desired from the determined DNA sequences or on the basis of the corresponding amino acid sequences.

DNA coding for acceptor framework sequences is widely available to those skilled in the art and can be readily synthesised on the basis of their known amino acid sequences.

10 Standard techniques of molecular biology may be used to prepare DNA sequences coding for the antibody molecule of the present invention. Desired DNA sequences may be synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate.

Any suitable host cell/vector system may be used for expression of the DNA  
15 sequences encoding the antibody molecule of the present invention. Bacterial, for example *E. coli*, and other microbial systems may be used, in part, for expression of antibody fragments such as Fab and F(ab')<sub>2</sub> fragments, and especially Fv fragments and single chain antibody fragments, for example, single chain Fvs. Eukaryotic, e.g. mammalian, host cell expression systems may be used for production of larger antibody molecules, including  
20 complete antibody molecules. Suitable mammalian host cells include CHO, myeloma or hybridoma cells.

The present invention also provides a process for the production of an antibody molecule according to the present invention comprising culturing a host cell comprising a vector of the present invention under conditions suitable for leading to expression of  
25 protein from DNA encoding the antibody molecule of the present invention, and isolating the antibody molecule.

Preferably the process for the production of the antibody molecule of the present invention comprises culturing *E. coli* comprising an *E. coli* expression vector comprising the DNA sequence of the present invention under conditions suitable for leading to  
30 expression of protein from the DNA sequence and isolating the antibody molecule. The antibody molecule may be secreted from the cell or targeted to the periplasm by suitable signal sequences. Alternatively, the antibody molecules may accumulate within the cell's cytoplasm. Preferably the antibody molecule is targeted to the periplasm. Depending on

the antibody molecule being produced and the process used, it is desirable to allow the antibody molecules to refold and adopt a functional conformation. Procedures for allowing antibody molecules to refold are well known to those skilled in the art.

The antibody molecule may comprise only a heavy or light chain polypeptide, in which case only a heavy chain or light chain polypeptide coding sequence needs to be used to transfect the host cells. For production of products comprising both heavy and light chains, the cell line may be transfected with two vectors, a first vector encoding a light chain polypeptide and a second vector encoding a heavy chain polypeptide. Alternatively, a single vector may be used, the vector including sequences encoding light chain and heavy chain polypeptides.

The present invention also provides a therapeutic or diagnostic composition comprising an antibody molecule of the present invention in combination with a pharmaceutically acceptable excipient, diluent or carrier.

The present invention also provides a process for preparation of a therapeutic or diagnostic composition comprising admixing the antibody molecule of the present invention together with a pharmaceutically acceptable excipient, diluent or carrier.

The antibody molecule may be the sole active ingredient in the therapeutic or diagnostic composition or may be accompanied by other active ingredients including other antibody ingredients, for example anti-T cell, anti-IFN $\gamma$  or anti-LPS antibodies, or non-antibody ingredients such as xanthines.

The pharmaceutical compositions should preferably comprise a therapeutically effective amount of the antibody of the invention. The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent needed to treat, ameliorate or prevent a targeted disease or condition, or to exhibit a detectable therapeutic or preventative effect. For any antibody, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually in rodents, rabbits, dogs, pigs or primates. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

The precise effective amount for a human subject will depend upon the severity of the disease state, the general health of the subject, the age, weight and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities and tolerance/response to therapy. This amount can be determined by routine experimentation

and is within the judgement of the clinician. Generally, an effective dose will be from 0.01 mg/kg to 50 mg/kg, preferably 0.1 mg/kg to 20 mg/kg, more preferably about 15 mg/kg. As shown in the Examples below, doses of 1, 5 and 20 mg/kg have been used to treat patients suffering from rheumatoid arthritis.

- 5           Compositions may be administered individually to a patient or may be administered in combination with other agents, drugs or hormones.

The dose at which the antibody molecule of the present invention is administered depends on the nature of the condition to be treated, the degree to which the level of TNF $\alpha$  to be neutralised is, or is expected to be, raised above a desirable level, and on whether the  
10   antibody molecule is being used prophylactically or to treat an existing condition.

Thus, for example, where the product is for treatment or prophylaxis of a chronic inflammatory disease, such as rheumatoid arthritis, suitable doses of the antibody molecule of the present invention lie in the range of between 0.5 and 50 mg/kg, more preferably between 1 and 20 mg/kg and most preferably about 15 mg/kg. The frequency of dose will  
15   depend on the half-life of the antibody molecule and the duration of its effect.

If the antibody molecule has a short half-life (e.g. 2 to 10 hours) it may be necessary to give one or more doses per day. Alternatively, if the antibody molecule has a long half life (e.g. 2 to 15 days) it may only be necessary to give a dosage once per day, per week or even once every 1 or 2 months.

- 20           A pharmaceutical composition may also contain a pharmaceutically acceptable carrier for administration of the antibody. The carrier should not itself induce the production of antibodies harmful to the individual receiving the composition and should not be toxic. Suitable carriers may be large, slowly metabolised macromolecules such as proteins, polypeptides, liposomes, polysaccharides, polylactic acids, polyglycolic acids,  
25   polymeric amino acids, amino acid copolymers and inactive virus particles.

Pharmaceutically acceptable salts can be used, for example mineral acid salts, such as hydrochlorides, hydrobromides, phosphates and sulphates, or salts of organic acids, such as acetates, propionates, malonates and benzoates.

- Pharmaceutically acceptable carriers in therapeutic compositions may additionally  
30   contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents or pH buffering substances, may be present in such compositions. Such carriers enable the pharmaceutical compositions to be

formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries and suspensions, for ingestion by the patient.

Preferred forms for administration include forms suitable for parenteral administration, e.g. by injection or infusion, for example by bolus injection or continuous  
5 infusion. Where the product is for injection or infusion, it may take the form of a suspension, solution or emulsion in an oily or aqueous vehicle and it may contain formulatory agents, such as suspending, preservative, stabilising and/or dispersing agents. Alternatively, the antibody molecule may be in dry form, for reconstitution before use with an appropriate sterile liquid.

10 Once formulated, the compositions of the invention can be administered directly to the subject. The subjects to be treated can be animals. However, it is preferred that the compositions are adapted for administration to human subjects.

The pharmaceutical compositions of this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-  
15 arterial, intramedullary, intrathecal, intraventricular, transdermal, transcutaneous (for example, see WO98/20734), subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, intravaginal or rectal routes. Hyposprays may also be used to administer the pharmaceutical compositions of the invention. Typically, the therapeutic compositions may be prepared as injectables, either as liquid solutions or suspensions. Solid forms suitable  
20 for solution in, or suspension in, liquid vehicles prior to injection may also be prepared.

Direct delivery of the compositions will generally be accomplished by injection, subcutaneously, intraperitoneally, intravenously or intramuscularly, or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Dosage treatment may be a single dose schedule or a multiple dose schedule.

25 It will be appreciated that the active ingredient in the composition will be an antibody molecule. As such, it will be susceptible to degradation in the gastrointestinal tract. Thus, if the composition is to be administered by a route using the gastrointestinal tract, the composition will need to contain agents which protect the antibody from degradation but which release the antibody once it has been absorbed from the  
30 gastrointestinal tract.

A thorough discussion of pharmaceutically acceptable carriers is available in Remington's Pharmaceutical Sciences (Mack Publishing Company, N.J. 1991).



It is also envisaged that the antibody of the present invention will be administered by use of gene therapy. In order to achieve this, DNA sequences encoding the heavy and light chains of the antibody molecule under the control of appropriate DNA components are introduced into a patient such that the antibody chains are expressed from the DNA sequences and assembled *in situ*.

The present invention also provides the antibody molecule of the present invention for use in treating a disease mediated by TNF $\alpha$ .

The present invention further provides the use of the antibody molecule according to the present invention in the manufacture of a medicament for the treatment of a disease mediated by TNF $\alpha$ .

The antibody molecule of the present invention may be utilised in any therapy where it is desired to reduce the level of biologically active TNF $\alpha$  present in the human or animal body. The TNF $\alpha$  may be circulating in the body or present in an undesirably high level localised at a particular site in the body.

For example, elevated levels of TNF $\alpha$  are implicated in acute and chronic immune and immunoregulatory disorders, infections including septic, endotoxic and cardiovascular shock, inflammatory disorders, neurodegenerative diseases, malignant diseases and alcohol induced hepatitis. Details of the numerous disorders associated with elevated levels of TNF $\alpha$  are set out in US-A-5 919 452. The antibody molecule of the present invention may be utilised in the therapy of diseases mediated by TNF $\alpha$ . Particularly relevant diseases which may be treated by the antibody molecule of the present invention include sepsis, congestive heart failure, septic or endotoxic shock, cachexia, adult respiratory distress syndrome, AIDS, allergies, psoriasis, TB, inflammatory bone disorders, blood coagulation disorders, burns, rejection episodes following organ or tissue transplant, Crohn's disease and autoimmune diseases, such as thyroiditis and rheumatoid- and osteo-arthritis.

Additionally, the antibody molecule or composition may be used: to reduce side effects associated with TNF $\alpha$  generation during neoplastic therapy; to eliminate or reduce shock-related symptoms associated with the treatment or prevention of graft rejection by use of an anti-lymphocyte antibody; or for treating multi-organ failure.

The antibody molecule of the present invention is preferably used for treatment of rheumatoid- or osteo-arthritis.

The present invention also provides a method of treating human or animal subjects suffering from or at risk of a disorder mediated by TNF $\alpha$ , the method comprising

administering to the subject an effective amount of the antibody molecule of the present invention.

The antibody molecule of the present invention may also be used in diagnosis, for example in the *in vivo* diagnosis and imaging of disease states involving elevated levels of

5 TNF $\alpha$ .

The present invention also provides an antibody molecule comprising a hybrid CDR comprising a truncated donor CDR sequence wherein the missing portion of the truncated donor CDR is replaced by a different sequence and forms a functional CDR. The term "hybrid CDR" as used herein means a CDR comprising a donor CDR which has been

10 truncated at one or more positions, for example at one or both of its ends. The missing portion of the truncated donor CDR is replaced by a different sequence to form a complete and functional CDR. The hybrid CDR has at least one amino acid change compared to the complete donor CDR. The sequence replacing the truncated portion of the CDR can be any sequence. Preferably the non-donor part of the CDR sequence is from the antibody from

15 which the framework regions of the antibody molecule are derived, such as a germline antibody sequence.

It has been found that antibody molecules comprising a hybrid CDR retain substantially the same binding affinity as an antibody molecule comprising complete donor CDRs. The term "substantially the same binding affinity" as used herein means at least

20 70%, more preferably at least 85% and most preferably at least 95% of the binding affinity of the corresponding antibody molecule comprising complete donor CDRs. As noted above, in certain cases, the affinity of the antibody of the invention may be greater than that of the donor antibody. The use of a hybrid CDR provides the advantages of reducing the amount of foreign (i.e. donor) sequence present in the antibody molecule and may increase

25 the binding affinity of the antibody molecule compared to the corresponding antibody molecule comprising complete donor CDRs.

Any of the CDRs of the antibody molecule can be hybrid. Preferably CDR2 of the heavy chain is hybrid in the antibody molecule.

Preferably the truncation of the donor CDR is from 1 to 8 amino acids, more

30 preferably from 4 to 6 amino acids. It is further preferred that the truncation is made at the C-terminus of the CDR.

Depending on the sequence of the truncated portion of the CDR and the sequence of the different sequence replacing the missing portion, a number of amino acid changes may

be made. Preferably at least 2 amino acid changes are made, more preferably at least 3 amino acid changes are made and most preferably at least 4 amino acid changes are made.

A particular embodiment of this aspect of the invention is an antibody according to the first aspect of the invention wherein the second CDR in the heavy chain has the  
5 sequence given as SEQ ID NO:2. This has better affinity for its antigen than does the donor antibody from which part of the CDR is derived.

The present invention also provides a nucleic acid sequence which encodes the antibody molecule comprising a hybrid CDR of the present invention.

The present invention also provides an expression vector containing the nucleic acid  
10 sequence encoding the antibody molecule comprising a hybrid CDR of the present invention.

The present invention also provides a host cell transformed with the vector of the present invention.

The present invention also provides a process for the production of an antibody  
15 molecule comprising a hybrid CDR comprising culturing the host cell of the present invention and isolating the antibody molecule.

The present invention is further described by way of illustration only in the following examples which refer to the accompanying Figures, in which:

Figure 1 shows the framework regions of the human light chain subgroup 1  
20 compared to the framework regions of the hTNF40 light chain (SEQ ID NOS:83 to 90);

Figure 2 shows the framework regions of the human heavy chain subgroup 1 and subgroup 3 compared to the framework regions of the hTNF40 heavy chain (SEQ ID NOS:91 to 98 and 106 to 109);

Figure 3 shows the amino acid sequence of the CDRs of hTNF40 (SEQ ID NOS:1  
25 to 7), wherein CDR H2' is a hybrid CDR wherein the C-terminal six amino acids are from the H2 CDR sequence of a human subgroup 3 germline antibody and the amino acid changes to the sequence resulting from this hybridisation are underlined;

Figure 4 shows vector pMR15.1;

Figure 5 shows vector pMR14;

30 Figure 6 shows the nucleotide and predicted amino acid sequence of the murine hTNF40V1 (SEQ ID NO: 99);

Figure 7 shows the nucleotide and predicted amino acid sequence of the murine hTNF40Vh (SEQ ID NO:100);

Figure 8 shows the nucleotide and predicted amino acid sequence of hTNF40-gL1 (SEQ ID NO:8);

Figure 9 shows the nucleotide and predicted amino acid sequence of hTNF40-gL2 (SEQ ID NO:9);

5 Figure 10 shows the nucleotide and predicted amino acid sequence of gh1hTNF40.4 (SEQ ID NO:10);

Figure 11 shows the nucleotide and predicted amino acid sequence of gh3hTNF40.4 (SEQ ID NO:11);

Figure 12 shows vector CTIL5-gL6;

10 Figure 13 shows the structure of a compound called CDP870 comprising a modified Fab fragment derived from antibody hTNF40 covalently linked via a cysteine residue to a lysyl-maleimide linker wherein each amino group on the lysyl residue has covalently attached to it a methoxy PEG residue wherein n is about 420;

Figure 14 shows vector pTTQ9;

15 Figure 15 shows the sequence of the OmpA oligonucleotide adapter (SEQ ID NO:101);

Figure 16 shows vector pACYC184;

Figure 17 shows vector pTTO-1;

Figure 18 shows vector pTTO-2;

20 Figure 19 shows vector pDNAbEng-G1;

Figure 20 shows the oligonucleotide cassettes encoding different intergenic sequences for *E. coli* modified Fab expression (SEQ ID NOS:102 to 105);

Figure 21 shows periplasmic modified Fab accumulation of IGS variants;

Figure 22 shows vector pTTO(CDP870);

25 Figure 23 shows the disease activity score (DAS) in patients treated with different doses of CDP870 and placebo. Median and IQ ranges are presented for the per-protocol population with last observation carried forward. Small squares indicate placebo, diamonds indicate 1 mg/kg, triangles indicate 5 mg/kg and large squares indicate 20 mg/kg;

30 Figure 24 shows the tender joint count, swollen joint count, pain score, assessor's global assessment of disease activity, modified health assessment questionnaire (HAQ), C reactive protein (CRP) and erythrocyte sedimentation rate (ESR) in patients treated with different doses of CDP870 and placebo. Median and IQ range are presented for the per-

protocol population with last observation carried forward. Small squares indicate placebo, diamonds indicate 1 mg/kg, triangles indicate 5 mg/kg and large squares indicate 20 mg/kg.

## **EXAMPLES**

5

### **Gene Cloning and Expression of a Chimeric hTNF40 Antibody Molecule**

#### **RNA Preparation from hTNF40 Hybridoma Cells**

Total RNA was prepared from  $3 \times 10^7$  hTNF40 hybridoma cells as described below.

10 Cells were washed in physiological saline and dissolved in RNazol (0.2 ml per  $10^6$  cells). Chloroform (0.2 ml per 2 ml homogenate) was added, the mixture shaken vigorously for 15 seconds and then left on ice for 15 minutes. The resulting aqueous and organic phases were separated by centrifugation for 15 minutes in an Eppendorf centrifuge and RNA was precipitated from the aqueous phase by the addition of an equal volume of isopropanol.

15 After 15 minutes on ice, the RNA was pelleted by centrifugation, washed with 70% ethanol, dried and dissolved in sterile, RNase free water. The yield of RNA was 400 µg.

#### **PCR Cloning of hTNF40 Vh and VI**

cDNA sequences coding for the variable domains of hTNF40 heavy and light chains

20 were synthesised using reverse transcriptase to produce single stranded cDNA copies of the mRNA present in the total RNA, followed by Polymerase Chain Reaction (PCR) on the cDNAs with specific oligonucleotide primers.

##### **a) cDNA Synthesis**

cDNA was synthesised in a 20 µl reaction volume containing the following

25 reagents: 50mM Tris-HCl pH 8.3, 75 mM KCl, 10 mM dithiothreitol, 3 mM  $MgCl_2$ , 0.5 mM each deoxyribonucleoside triphosphate, 20 units RNAsin, 75 ng random hexanucleotide primer, 2 µg hTNF40 RNA and 200 units Moloney Murine Leukemia Virus reverse transcriptase. After incubation at 42°C for 60 minutes, the reaction was terminated by heating at 95°C for 5 minutes.

##### **30 b) PCR**

Aliquots of the cDNA were subjected to PCR using combinations of primers specific for the heavy and light chains. The nucleotide sequences of the 5' primers for the heavy and light chains are shown in Tables 1 and 2 respectively. These sequences all contain, in order, a

restriction site starting 7 nucleotides from their 5' ends, the sequence GCCGCCACC (SEQ ID NO:12), to allow optimal translation of the resulting mRNAs, an initiation codon and 20-30 nucleotides based on the leader peptide sequences of known mouse antibodies (Kabat *et al.*, Sequences of proteins of immunological interest, 5<sup>th</sup> Edition, 1991, U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health).

The 3' primers are shown in Table 3. The light chain primer spans the J-C junction of the antibody and contains a restriction site for the enzyme Sp1I to facilitate cloning of the V1 PCR fragment. The heavy chain 3' primers are a mixture designed to span the J-C junction of the antibody. The 3' primer includes an ApaI restriction site to facilitate cloning. The 3' region of the primers contains a mixed sequence based on those found in known mouse antibodies (Kabat *et al.*, 1991, *supra*).

The combinations of primers described above enable the PCR products for Vh and V1 to be cloned directly into an appropriate expression vector (see below) to produce chimeric (mouse-human) heavy and light chains and for these genes to be expressed in mammalian cells to produce chimeric antibodies of the desired isotype.

Incubations (100 µl) for the PCR were set up as follows. Each reaction contained 10 mM Tris-HCl pH 8.3, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, 0.01% w/v gelatin, 0.25 mM each deoxyribonucleoside triphosphate, 10 pmoles 5' primer mix (Table 4), 10 pmoles 3' primer (CL12 (light chain) or R2155 (heavy chain) (Table 3)), 1 µl cDNA and 1 unit Taq polymerase. Reactions were incubated at 95°C for 5 minutes and then cycled through 94°C for 1 minute, 55°C for 1 minute and 72°C for 1 minute. After 30 cycles, aliquots of each reaction were analysed by electrophoresis on an agarose gel. Light chain reactions containing 5' primer mixes from light chain pools 1, 2 and 7 produced bands with sizes consistent with full length V1 fragments while the reaction from heavy chain reaction pool 3 produced a fragment with a size expected of a Vh gene. The band produced by the light chain pool 1 primers was not followed up as previous results had shown that this band corresponds to a light chain pseudogene produced by the hybridoma cell. The band produced by the light chain pool 7 primers was weaker than the band from the pool 2 primers and therefore was not followed up. Only the band from light chain reaction pool 2, which was the strongest band, was followed up.

### c) Molecular Cloning of the PCR Fragments

The DNA fragments produced in the light chain reaction pool 2 were digested with the enzymes BstBI and Sp1I, concentrated by ethanol precipitation, electrophoresed on a

1.4% agarose gel and DNA bands in the range of 400 base pairs recovered. These were cloned by ligation into the vector pMR15.1 (Figure 4) that had been restricted with BstBI and SphI. After ligation, mixtures were transformed into *E. coli* LM 1035 and plasmids from the resulting bacterial colonies screened for inserts by digestion with BstBI and SphI.

- 5 Representatives with inserts from each ligation were analysed further by nucleotide sequencing.

In a similar manner, the DNA fragments produced in heavy chain reaction pool 3 were digested with HindIII and ApaI and cloned into the vector pMR14 (Figure 5) that had been restricted with HindIII and ApaI. Again, representative plasmids containing inserts  
10 were analysed by nucleotide sequencing.

#### d) Nucleotide Sequence Analysis

Plasmid DNA from a number of isolates containing Vh inserts was sequenced using the primers R1053 (see Table 5) (which primes in the 3' region of the HCMV promoter in pMR14) and R720 (see Table 5) (which primes in the 5' region of human C – gamma 4 and  
15 allows sequencing through the DNA insert on pMR14). It was found that the nucleotide sequences of the Vh insert in a number of clones were identical, except for differences in the signal peptide and J regions. This indicated that the clones examined are independent isolates arising from the use of different primers from the mixture of oligonucleotides during the PCR stage. The determined nucleotide sequence and predicted amino acid  
20 sequence of the variable domain of the heavy chain of antibody hTNF40 (hTNF40Vh) are given in Figure 7 (SEQ ID NO:100).

To analyse the light chain clones, the sequence derived from priming with R1053 (see Table 5) and R684 (SEQ ID NO:62) (which primes in the 5' region of human C-kappa and allows sequencing through the DNA insert on pMR15.1) was examined. The  
25 nucleotide sequence and predicted amino acid sequence of the V<sub>L</sub> genes arising from reactions in pool 2 were similarly analysed. Again it was found that the nucleotide sequences of the V<sub>L</sub> insert in a number of clones were identical, except for differences in the signal peptide and J regions, indicating that the clones examined were independent isolates arising from the use of different primers from the mixture of oligonucleotides used during  
30 the PCR stage. The determined nucleotide sequence and predicted amino acid sequence of the variable domain of the light chain of antibody hTNF40 (hTNF40V<sub>L</sub>) are given in Figure 6 (SEQ ID NO:99).

**TABLE 1**

Oligonucleotide primers for the 5' region of mouse heavy chains.

- 5 CH1 : 5'ATGAAATGCAGCTGGGTCAT(G,C)TTCTT3' (SEQ ID NO:13)  
 CH2 : 5'ATGGGATGGAGCT(A,G)TATCAT(C,G)(C,T)TCTT3' (SEQ ID NO:14)  
 CH3 : 5'ATGAAG(A,T)TGTGGTTAAACTGGGTTTT3' (SEQ ID NO:15)  
 CH4 : 5'ATG(G,A)ACTTTGGG(T,C)TCAGCTTG(G,A)T3' (SEQ ID NO:16)  
 CH5 : 5'ATGGACTCCAGGCTCAATTTAGTTTT3' (SEQ ID NO:17)  
 10 CH6 : 5'ATGGCTGTC(C,T)T(G,A)G(G,C)GCT(G,A)CTCTTCTG3' (SEQ ID NO:18)  
 CH7 : 5'ATGG(G,A)ATGGAGC(G,T)GG(G,A)TCTTT(A,C)TCTT3' (SEQ ID NO:19)  
 CH8 : 5'ATGAGAGTGCTGATTCTTTTGTG3' (SEQ ID NO:20)  
 CH9 : 5'ATGG(C,A)TTGGGTGTGGA(A,C)CTTGCTATT3' (SEQ ID NO:21)  
 CH10 : 5'ATGGGCAGACTTACATTCTCATTCT3' (SEQ ID NO:22)  
 15 CH11 : 5'ATGGATTTTGGGCTGATTTTTTTTATTG3' (SEQ ID NO:23)  
 CH12 : 5'ATGATGGTGTAAAGTCTTCTGTACCT3' (SEQ ID NO:24)

Each of the above primers has the sequence 5'GCGCGCAAGCTTGCCGCCACCT3' (SEQ ID NO:25) added to its 5' end.

20

**TABLE 2**

Oligonucleotide primers for the 5' region of mouse light chains.

- CL1 : 5'ATGAAGTTGCCTGTTAGGCTGTTGGTGCT3' (SEQ ID NO:26)  
 25 CL2 : 5'ATGGAG(T,A)CAGACACACTCCTG(T,C)TATGGGT3' (SEQ ID NO:27)  
 CL3 : 5'ATGAGTGTGCTCACTCAGGTCCT3' (SEQ ID NO:28)  
 CL4 : 5'ATGAGG(G,A)CCCCTGCTCAG(A,T)TT(C,T)TTGG3' (SEQ ID NO:29)  
 CL5 : 5'ATGGATTT(T,A)CAGGTGCAGATT(T,A)TCAGCTT3' (SEQ ID NO:30)  
 CL5A : 5'ATGGATTT(T,A)CA(A,G)GTGCAGATT(T,A)TCAGCTT3' (SEQ ID NO:31)  
 30 CL6 : 5'ATGAGGT(T,G)C(T,C)(T,C)TG(T,C)T(G,C)AG(T,C)T(T,C)CTG(A,G)G3' (SEQ ID NO:32)  
 CL7 : 5'ATGGGC(T,A)TCAAGATGGAGTCACA3' (SEQ ID NO:33)



- CL8 : 5'ATGTGGGGA(T,C)CT(G,T)TTT(T,C)C(A,C)(A,C)TTTTTCAAT3'(SEQ ID NO:34)
- CL9 : 5'ATGGT(G,A)TCC(T,A)CA(G,C)CTCAGTTCCTT3' (SEQ ID NO:35)
- CL10 : 5'ATGTATATATGTTTGTGTCTATTTC3' (SEQ ID NO:36)
- 5 CL11 : 5'ATGGAAGCCCCAGCTCAGCTTCTCTT3'(SEQ ID NO:37)
- CL12A : 5'ATG(A,G)AGT(T,C)(A,T)CAGACCCAGGTCTT(T,C)(A,G)T3' (SEQ ID NO:38)
- CL12B : 5'ATGGAGACACATTCTCAGGTCTTTGT3' (SEQ ID NO:39)
- CL13 : 5'ATGGATTACAGGCCCCAGGTTCCTTAT3' (SEQ ID NO:40)
- 10 CL14 : 5'ATGATGAGTCCTGCCCAGTTCCTGTT3' (SEQ ID NO:41)
- CL15 : 5'ATGAATTTGCCTGTTTCATCTCTTGGTGCT3' (SEQ ID NO:42)
- CL16 : 5'ATGGATTTTCAATTGGTCCTCATCTCCTT3' (SEQ ID NO:43)
- CL17A : 5'ATGAGGTGCCTA(A,G)CT(C,G)AGTTCCTG(A,G)G3' (SEQ ID NO:44)
- CL17B : 5'ATGAAGTACTCTGCTCAGTTTCTAGG3' (SEQ ID NO:45)
- 15 CL17C : 5'ATGAGGCATTCTCTTCAATTCTTGGG3' (SEQ ID NO:46)

Each of the above primers has the sequence 5'GGACTGTTCTGAAGCCGCCACCC3' (SEQ ID NO:47) added to its 5' end.

## 20 TABLE 3

Oligonucleotide primers for the 3' ends of mouse Vh and Vl genes.

Light chain (CL12) :

5'GGATACAGTTGGTGCAGCATCCGTACGTTT3' (SEQ ID NO:48)

25

Heavy chain ( R2155 ) :

5'GCAGATGGGCCCTTCGTTGAGGCTG(A,C)(A,G)GAGAC(G,T,A)GTGA3'  
(SEQ ID NO:49)

**TABLE 4**

## a) 5' Primer mixtures for light chain PCR reactions

- 
- pool 1 : CL2.  
 5 pool 2 : CL7.  
 pool 3 : CL13.  
 pool 4 : CL6.  
 pool 5 : CL5A, CL9, CL17A.  
 pool 6 : CL8.  
 10 pool 7 : CL12A.  
 pool 8 : CL1, CL3, CL4, CL5, CL10, CL11, CL2B, CL14, CL15, CL16, CL17B, CL17C

## b) 5' Primer mixtures for heavy chain PCR reactions

- 15 pool 1 : CH1, CH2, CH3, CH4.  
 pool 2 : CH5, CH6, CH7, CH8.  
 pool 3 : CH9, CH10, CH11, CH12.

**Table 5**

## 20 Primers used in nucleotide sequence analysis

- R1053 : 5'GCTGACAGACTAACAGACTGTTCC3' (SEQ ID NO:50)  
 R720 : 5'GCTCTCGGAGGTGCTCCT3' (SEQ ID NO:51)

25 **Evaluation of Activities of Chimeric Genes**

The activities of the chimeric genes were evaluated by expressing them in mammalian cells and purifying and quantitating the newly synthesised antibodies. The methodology for this is described below, followed by a description of the biochemical and cell based assays used for the biological characterisation of the antibodies.

30 a) **Production of Chimeric hTNF40 Antibody Molecule**

Chimeric antibody for biological evaluation was produced by transient expression of the appropriate heavy and light chain pairs after co-transfection into Chinese Hamster Ovary (CHO) cells using calcium phosphate precipitation.

On the day prior to transfection, semi-confluent flasks of CHO-L761 cells were trypsinised, the cells counted and T75 flasks set up each with  $10^7$  cells.

On the next day, the culture medium was changed 3 hours before transfection. For transfection, the calcium phosphate precipitate was prepared by mixing 1.25 ml of 0.25 M  
5  $\text{CaCl}_2$  containing 50  $\mu\text{g}$  of each of heavy and light chain expression vectors with 1.25 ml of  
2 x HBS (16.36 g NaCl, 11.0 g HEPES and 0.4 g  $\text{Na}_2\text{HPO}_4$  in 1 litre water with the pH  
adjusted to 7.1 with NaOH) and adding immediately into the medium of the cells. After 3  
hours at  $37^\circ\text{C}$  in a  $\text{CO}_2$  incubator, the medium and precipitate were removed and the cells  
shocked by the addition of 15 ml 15% glycerol in phosphate buffered saline (PBS) for 1  
10 minute. The glycerol was removed, the cells washed once with PBS and incubated for 48-  
96 hours in 25 ml medium containing 10 mM sodium butyrate. Antibody could be purified  
from the culture medium by binding to and elution from protein A-Sepharose.

#### b) ELISA

For the ELISA, Nunc ELISA plates were coated overnight at  $4^\circ\text{C}$  with a  $\text{F(ab)}_2$   
15 fragment of a polyclonal goat anti-human Fc fragment specific antibody (Jackson  
ImmunoResearch, code 109-006-098) at 5  $\mu\text{g/ml}$  in coating buffer (15 mM sodium  
carbonate, 35 mM sodium hydrogen carbonate, pH 6.9). Uncoated antibody was removed  
by washing 5 times with distilled water. Samples and purified standards to be quantitated  
were diluted to approximately 1  $\mu\text{g/ml}$  in conjugate buffer (0.1 M Tris-HCl, pH 7.0, 0.1 M  
20 NaCl, 0.2% v/v Tween 20, 0.2% w/v Hammersten casein). The samples were titrated in  
the microtitre wells in 2-fold dilutions to give a final volume of 0.1 ml in each well and the  
plates incubated at room temperature for 1 hour with shaking. After the first incubation  
step the plates were washed 10 times with distilled water and then incubated for 1 hour as  
before with 0.1 ml of a mouse monoclonal anti-human kappa (clone GD12) peroxidase  
25 conjugated antibody (The Binding Site, code MP135) at a dilution of 1 in 700 in conjugate  
buffer. The plate was washed again and substrate solution (0.1 ml) added to each well.  
Substrate solution contained 150  $\mu\text{l}$  N,N,N,N-tetramethylbenzidine (10 mg/ml in DMSO),  
150  $\mu\text{l}$  hydrogen peroxide (30% solution) in 10 ml 0.1 M sodium acetate/sodium citrate, pH  
6.0. The plate was developed for 5-10 minutes until the absorbance at 630 nm was  
30 approximately 1.0 for the top standard. Absorbance at 630 nm was measured using a plate  
reader and the concentration of the sample determined by comparing the titration curves  
with those of the standard.

c) **Determination of Affinity constants by BiaCore analysis.**

The binding interaction between hTNF40 and human TNF was investigated using BIA technology. An affinity purified goat polyclonal antibody, directed against the constant region of hTNF40, was immobilised on the dextran polymer sensor chip surface using standard NHS/EDC chemistry. Relatively low levels (200-500 RU) of hTNF40 were captured to ensure mass transport effects were minimised. Human TNF at different concentrations was passed over the captured hTNF40 to allow assessment of the association kinetics. Following the injection of ligand, buffer was passed over the surface so that the dissociation could be measured. The association and dissociation rate constants for the interaction between solid phase hTNF40 and human TNF were calculated, and a  $K_D$  value was derived.

**EXAMPLE 1**

**CDR-Grafting of hTNF40**

The molecular cloning of genes for the variable regions of the heavy and light chains of the hTNF40 antibody and their use to produce chimeric (mouse-human) hTNF40 antibodies has been described above. The nucleotide and amino acid sequences of the murine hTNF40 V<sub>L</sub> and V<sub>H</sub> are shown in Figures 6 and 7 (SEQ ID NOS:99 and 100), respectively. This example describes the CDR-grafting of the hTNF40 antibody.

**CDR-Grafting of hTNF40 Light Chain**

Alignment of the framework regions of hTNF40 light chain with those of the four human light chain subgroups (Kabat *et al.*, 1991, *supra*) revealed that hTNF40 was most homologous to antibodies in human light chain subgroup 1. Consequently, for constructing the CDR-grafted light chain, the framework regions chosen corresponded to those of the human group 1 consensus sequence.

A comparison of the amino acid sequences of the framework regions of murine hTNF40 and the consensus human group 1 light chains is given in Figure 1 and shows that there are 22 differences (underlined) between the two sequences. Analysis of the contribution that any of these framework differences might have on antigen binding identified 2 residues for investigation; these are at positions 46 and 60. Based on this analysis, two versions of the CDR-grafted light chain were constructed. In the first of these, hTNF40-gL1 (SEQ ID NO:8), residues 46 and 60 are derived from the hTNF40 light

chain while in the second, hTNF40-gL2 (SEQ ID NO:9), all residues are human consensus except residue number 60 which is from the hTNF40 light chain.

#### **Construction of CDR-Grafted Light Chain hTNF40-gL1.**

5        The construction of hTNF40-gL1 is given below in detail. The following overlapping oligonucleotides (P7982-P7986) were used in the Polymerase Chain Reactions (PCR) to assemble a truncated grafted light chain. The assembled fragment lacks the antibody leader sequence and the first 17 amino acids of framework 1.

10    oligo 1 P7982:

5' GAATTCAGGGTCACCATCACTTGTAAGCCAGTCAGAACGTAGGTACTAAC  
GTAGCCTGGTATCAGCAA3' (SEQ ID NO:52)

oligo 2 P7983:

15    5' ATAGAGGAAAGAGGCACTGTAGATGAGGGCTTTGGGGCTTTACCTGGTTT  
TTGCTGATACCAGGCTACGT3' (SEQ ID NO:53)

oligo 3 P7984:

20    5' TACAGTGCCTCTTTCCTCTATAGTGGTGTACCATACAGGTTTCAGCGGATCCG  
GTAGTGGTACTGATTTCAC3' (SEQ ID NO:54)

oligo 4 P7985

5'GACAGTAATAAGTGGCGAAATCTTCTGGCTGGAGGCTACTGATCGTGAGGGT  
GAAATCAGTACCACTACCG3' (SEQ ID NO:55)

25

oligo 5 P7986:

5'ATTTCCGCACTTATTACTGTCAACAGTATAACATCTACCCACTCACATTCGGT  
CAGGGTACTAAAGTAGAAATCAAACGTACGGAATTC3' (SEQ ID NO:56)

30    Fwd P7981:

5'GAATTCAGGGTCACCATCACTTGTAAGCC3' (SEQ ID NO:57)

Bwd P7980

5'GAATTCCGTACGTTTGATTTCTACTTTAGT3' (SEQ ID NO:58),

- A PCR reaction, 100  $\mu$ l, was set up containing, 10 mM Tris-HCl pH 8.3, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, 0.01% w/v gelatin, 0.25 mM each deoxyribonucleoside triphosphate, 2 pmoles of P7982, P7983, P7984, P7985, P7986, 10 pmoles of P7980, P7981 and 1 unit of Taq polymerase. Reactions were cycled through 94°C for 1 minute, 55°C for 1 minute and 72°C for 1 minute. After 30 cycles, each reaction was analysed by electrophoresis on an agarose gel and the PCR fragment excised from the gel and recovered using a Mermaid Kit.
- The recovered fragment was restricted with the enzymes BstEII and SphI in the appropriate buffer. The resulting product was finally electrophoresed on an agarose gel and the 270 base pair DNA fragment recovered from a gel slice and ligated into vector CTIL5-gL6 (Figure 12), that had previously been digested with the same enzymes. The above vector provides the missing antibody leader sequence and the first 17 amino acids of framework 1.
- The ligation mixture was used to transform E. coli strain LM1035 and resulting colonies analysed by PCR, restriction enzyme digests and nucleotide sequencing. The nucleotide and amino acid sequence of the VI region of hTNF40-gL1 is shown in Figure 8 (SEQ ID NO:8).

## 20 Construction of CDR-Grafted Light Chain hTNF40-gL2.

hTNF40-gL2 (SEQ ID NO:9) was constructed using PCR. The following oligonucleotides were used to introduce the amino acid changes:

- R1053: 5'GCTGACAGACTAACAGACTGTTCC3' (SEQ ID NO:59)
- R5350: 5'TCTAGATGGCACACCATCTGCTAAGTTTGATGCAGCATAGAT  
CAGGAGCTTAGGAGC3' (SEQ ID NO:60)
- R5349: 5'GCAGATGGTGTGCCATCTAGATTTCAGTGGCAGTGGATCA  
GGCACAGACTTTACCCTAAC3' (SEQ ID NO:61)
- R684: 5'TTCAACTGCTCATCAGAT3' (SEQ ID NO:62)

Two reactions, each 20  $\mu$ l, were set up each containing 10 mM Tris-HCl pH 8.3, 1.5 mM  $MgCl_2$ , 50 mM KCl, 0.01% w/v gelatin, 0.25 mM each deoxyribonucleoside triphosphate, 0.1  $\mu$ g hTNF40-gL1, 6 pmoles of R1053/R5350 or R5349/R684 and 0.25 units Taq polymerase. Reactions were cycled through 94°C for 1 minute, 55°C for 1 minute and 72°C for 1 minute. After 30 cycles, each reaction was analysed by electrophoresis on an agarose gel and the PCR fragments excised from the gel and recovered using a Mermaid Kit.

Aliquots of these were then subjected to a second round of PCR. The reaction, 100  $\mu$ l, contained 10 mM Tris-HCl pH 8.3, 1.5 mM  $MgCl_2$ , 50 mM KCl, 0.01% w/v gelatin, 1/5 of each of the PCR fragments from the first set of reactions, 30 pmoles of R1053 and R684 and 2.5 units Taq polymerase. Reaction temperatures were as above. After the PCR, the mixture was extracted with phenol/chloroform and then with chloroform and precipitated with ethanol. The ethanol precipitate was recovered by centrifugation, dissolved in the appropriate buffer and restricted with the enzymes BstEII and SphI. The resulting product was finally electrophoresed on an agarose gel and the 270 base pair DNA fragment recovered from a gel slice and ligated into the vector pMR15.1 (Figure 4) that had previously been digested with the same enzymes.

The ligation mixture was used to transform E. coli LM1035 and resulting colonies analysed by PCR, restriction enzyme digests and nucleotide sequencing. The nucleotide and amino acid sequence of the VI region of hTNF40-gL2 is shown in Figure 9 (SEQ ID NO:9).

### CDR-Grafting of hTNF40 Heavy Chain

CDR-grafting of hTNF40 heavy chain was accomplished using the same strategy as described for the light chain. hTNF40 heavy chain was found to be most homologous to human heavy chains belonging to subgroup 1 and therefore the consensus sequence of the human subgroup 1 frameworks was chosen to accept the hTNF40 heavy chain CDRs.

To investigate the requirement of a homologous human framework to act as an acceptor framework for CDR grafting, a second framework, human group 3, was selected to humanise hTNF40 heavy chain.

A comparison of hTNF40 with the two different frameworks region is shown in Figure 2 where it can be seen that hTNF40 differs from the human subgroup 1 consensus at 32 positions (underlined) and differs from the human subgroup 3 consensus at 40 positions

(underlined). After analysis of the contribution that any of these might make to antigen binding, residues 28, 38, 46, 67, 69 and 71 were retained as donor in the CDR-grafted heavy chain gh1hTNF40.1, using the group 1 framework. Residues 27, 28, 30, 48, 49, 69, 71, 73, 76 and 78 were retained as donor in the CDR-grafted heavy chain, gh3hTNF40.4 using the group 3 framework. Residues 28, 69 and 71 were retained as donor in the CDR-grafted heavy chain, gh1hTNF40.4 using the group 1 framework.

#### Construction of CDR-Grafted Heavy Chain gh1hTNF40.4

gh1hTNF40.4 (SEQ ID NO:10) was assembled by subjecting overlapping oligonucleotides to PCR in the presence of the appropriate primers. The following oligonucleotides were used in the PCR:

##### Group 1 graft

oligo 1 P7989:

5'GAAGCACCAGGCTTCTTAACCTCTGCTCCTGACTGGACCAGCTGCACCTGAG  
AGTGCACGAATTC3' (SEQ ID NO:63)

oligo 2 P7990:

5'GGTTAAGAAGCCTGGTGCTTCCGTCAAAGTTTCGTGTAAGGCCTCAGGCTAC  
GTGTTACAGACTATGGTA3' (SEQ ID NO:64)

oligo 3 P7991:

5'CCAACCCATCCATTTTCAGGCCTTGTCCCGGGGCCTGCTTGACCCAATTCATAC  
CATAGTCTGTGAACACGT3' (SEQ ID NO:65)

oligo 4 P7995:

5'GGCCTGAAATGGATGGGTTGGATTAATACTTACATTGGAGAGCCTATTTATGT  
TGACGACTTCAAGGGCAGATTCACGTTC3' (SEQ ID NO:66)

oligo 5 P7992:

5'CCATGTATGCAGTGCGTTGTGGAGGTGTCTAGAGTGAACGTGAATCTGCCCTT  
GAA3' (SEQ ID NO:67)



oligo 6 P7993:

5'CCACAAGCACTGCATACATGGAGCTGTCATCTCTGAGATCCGAGGACACCGC  
AGTGTACTAT3' (SEQ ID NO:68)

5

oligo 7 P7994:

5'GAATTCGGTACCCTGGCCCCAGTAGTCCATGGCATAAGATCTGTATCCTCTAG  
CACAATAGTACACTGCGGTGTCCTC3' (SEQ ID NO:69)

10 Fwd: P7988:

5'GAATTCGTGCACTCTCAGGTGCAGCTGGTC3' (SEQ ID NO:70)

Bwd P7987:

5'GAATTCGGTACCCTGGCCCCAGTAGTCCAT3' (SEQ ID NO:71)

15

The assembly reaction, 100 µl, contained 10 mM Tris-HCl pH 8.3, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, 0.01% w/v gelatin, 0.25 mM each deoxyribonucleoside triphosphate, 2 pmole of each of p7989, p7990, p7991, p7995, p7992, p7993 and p7994, 10 pmoles of each of p7988 and p7987 and 1 unit Taq polymerase. Reactions were cycled through 94°C for 1  
20 minute, 55°C for 1 minute and 72°C for 1 minute. After 30 cycles, the reaction was extracted with phenol/chloroform (1/1), then with chloroform and precipitated with ethanol. After centrifugation, the DNA was dissolved in the appropriate restriction buffer and digested with ApaLI and KpnI. The resulting fragment was isolated from an agarose gel and ligated into pMR14 (Figure 5) that had previously been digested with the same  
25 enzymes. pMR14 contains the human gamma 4 heavy chain constant region when pMR14 is cleaved with ApaLI and KpnI, the cleaved vector is able to receive the digested DNA such that the 3' end of the digested DNA joins in reading frame to the 5' end of the sequence encoding the gamma 4 constant region. Therefore, the heavy chain expressed from this vector will be a gamma 4 isotype. The ligation mixture was used to transform *E.*  
30 *coli* LM1035 and resulting bacterial colonies screened by restriction digest and nucleotide sequence analysis. In this way, a plasmid was identified containing the correct sequence for gh1hTNF40.4 (Figure 10) (SEQ ID NO:10).

**Construction of CDR-Grafted Heavy Chain gh3hTNF40.4**

gh3hTNF40.4 (SEQ ID NO:11) was assembled by subjecting overlapping oligonucleotides to PCR in the presence of the appropriate primers. The following  
5 oligonucleotides were used in the PCR:

**Group 3 graft****oligo 1 P7999:**

10 5'GATCCGCCAGGCTGCACGAGACCGCCTCCTGACTCGACCAGCTGAACCTCAG  
AGTGCACGAATTC3' (SEQ ID NO:72)

**oligo 2 P8000:**

5'TCTCGTGCAGCCTGGCGGATCGCTGAGATTGTCCTGTGCTGCATCTGGTTACG  
15 TCTTCACAGACTATGGAA3' (SEQ ID NO:73)

**oligo 3 P8001**

5'CCAACCCATCCATTTCAGGCCCTTCCCGGGGCCTGCTTAACCCAATTCATTC  
CATAGTCTGTGAAGACGT3' (SEQ ID NO:74)  
20

**oligo 4 P7995:**

5'GGCCTGAAATGGATGGGTTGGATTAATACTTACATTGGAGAGCCTATTTATGT  
TGACGACTTCAAGGGCAGATTCACGTTTC3' (SEQ ID NO:66)

**25 oligo 5 P7997:**

5'GGAGGTATGCTGTTGACTTGGATGTGTCTAGAGAGAACGTGAATCTGCCCTT  
GAA3' (SEQ ID NO:75)

**oligo 6 P7998:**

30 5'CCAAGTCAACAGCATACCTCCAAATGAATAGCCTGAGAGCAGAGGACACCGC  
AGTGTACTAT3' (SEQ ID NO:76)

oligo 7 P7993:

5'GAATTCGGTACCCTGGCCCCAGTAGTCCATGGCATAAGATCTGTATCCTCTAG  
CACAATAGTACACTGCGGTGTCCTC3' (SEQ ID NO:77)

5 Fwd P7996:

5'GAATTCGTGCACTCTGAGGTTTCAGCTGGTC3' (SEQ ID NO:78)

Bwd P7987:

5'GAATTCGGTACCCTGGCCCCAGTAGTCCAT3' (SEQ ID NO:71)

10

The assembly reaction, 100 µl, contained 10 mM Tris-HCl pH 8.3, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, 0.01% w/v gelatin, 0.25 mM each deoxyribonucleoside triphosphate, 2 pmole of each of p7999, p8000, p8001, p7995, p7997, p7998 and p7993, 10 pmoles of each of p7996 and p7987 and 1 unit Taq polymerase. Reactions were cycled through 94°C for 1 minute, 55°C for 1 minute and 72°C for 1 minute. After 30 cycles, the reaction was extracted with phenol/chloroform (1/1), then with chloroform and precipitated with ethanol. After centrifugation, the DNA was dissolved in the appropriate restriction buffer and digested with ApaI and KpnI. The resulting fragment was isolated from an agarose gel and ligated into pMR14 (Figure 5) that had previously been digested with the same enzymes. pMR14 contained the human gamma 4 heavy chain constant region. When pMR14 is cleaved with ApaI and KpnI, the cleaved vector is able to receive the digested DNA such that the 3' end of the digested DNA joins in reading frame to the 5' end of the sequence encoding the gamma 4 constant region. Therefore, the heavy chain expressed from this vector will be a gamma 4 isotype. The ligation mixture was used to transform *E. coli* LM1035 and resulting bacterial colonies screened by restriction digestion and nucleotide sequence analysis. In this way, a plasmid was identified containing the correct sequence for gh3hTNF40.4 (SEQ ID NO:11) (Figure 11).

#### **Production of CDR-Grafted Modified Fab Fragment.**

30 A CDR-grafted, modified Fab fragment, based on antibody hTNF40, was constructed using the *E. coli* vector pTTO-1. The variable regions of antibody hTNF40 are sub-cloned into this vector and the intergenic sequence optimised to create pTTO(CDP870). The pTTO expression vector is designed to give rise to soluble,

periplasmic accumulation of recombinant proteins in *E. coli*. The main features of this plasmid are:

- (i) tetracycline resistance marker - antibiotic not inactivated by the product of resistance gene, hence selection for plasmid-containing cells is maintained;
- 5 (ii) low copy number - origin of replication derived from plasmid p15A, which is compatible with plasmids containing *colE1* derived replicons;
- (iii) strong, inducible *tac* promoter for transcription of cloned gene(s);
- (iv) *lacI<sup>q</sup>* gene - gives constitutive expression of the *lac* repressor protein, maintaining the *tac* promoter in the repressed state until induction with IPTG / allolactose;
- 10 (v) OmpA signal sequence - gives periplasmic secretion of cloned gene(s); and
- (vi) translational coupling of OmpA signal sequence to a short *lacZ* peptide, giving efficient initiation of translation.

The vector has been developed for expression of modified Fab fragments from a  
15 dicistronic message by the design of a method to select empirically the optimum intergenic sequence from a series of four purpose-built cassettes. The application of this in the construction of pTTO(CDP870) is described.

## Materials and Methods

20

### DNA techniques

Standard procedures were used for protocols including DNA restriction, agarose gel electrophoresis, ligation and transformation. Restriction enzymes and DNA modifying enzymes were obtained from New England Biolabs or Boehringer Mannheim, and were  
25 used according to the supplier's recommendations. DNA fragments were purified from agarose using the GeneClean protocol (BIO 101). Oligonucleotides were supplied by Oswel Oligonucleotide Service and were synthesized at the 40 nm scale. Plasmid DNA was isolated using Plasmid DNA Mini / Midi kits from Qiagen. PCR was performed using Perkin Elmer 'Amplitaq' as recommended. DNA sequencing was performed using the  
30 Applied Biosystems Taq cycle sequencing kit.

### Shake flask induction

*E. coli* W3110 cultures were grown in L-broth supplemented with tetracycline (7.5 µg/ml). For inductions, fresh overnight cultures (grown at 30°C) were diluted to OD<sub>600</sub> of

0.1 into 200 ml L-broth in a 2 L baffled flask and were grown at 30°C in an orbital incubator. At OD<sub>600</sub> of 0.5, IPTG was added to 200 µM. Samples (normalised for OD) were taken at intervals.

## 5 Periplasmic Extraction

Culture samples were chilled on ice (5 minutes) then cells were harvested by centrifugation. Following resuspension in extraction buffer (100 mM Tris.HCl, 10 mM EDTA, pH 7.4) samples were incubated overnight at 30°C, then clarified by centrifugation.

## 10 Assembly Assay

Modified Fab concentrations were determined by ELISA. Plates were coated at 4°C overnight with anti-human Fd 6045 (2 µg/ml in coating buffer, physiological saline, 100 µl per well). After washing, 100 µl of sample was loaded per well; purified A5B7 gamma-1 Fab', initially at 2 µg/ml, was used as a standard. Samples were serially diluted 2-fold across the plate in sample conjugate buffer (per litre: 6.05 g trisaminomethane; 2.92 g NaCl; 0.1 ml Tween-20; 1 ml casein (0.2%)); plates were incubated for 1 hour at room temperature, with agitation. Plates were washed and dried, then 100 µl of anti-human C-kappa (GD12)-peroxidase was added (diluted in sample conjugate buffer). Incubation was carried out at room temperature for 1 hour with agitation. Plates were washed and dried, then 100 µl of substrate solution was added (10 ml sodium acetate/citrate solution (0.1 M pH 6); 100 µl H<sub>2</sub>O<sub>2</sub> solution; 100 µl tetramethylbenzidine solution (10 mg/ml in dimethylsulphoxide)). Absorbance at 630 nm was read 4 - 6 minutes after substrate addition.

## 25 Construction of Plasmid pTTO-1

### (a) Replacement of the pTTQ9 Polylinker

Plasmid pTTQ9 was obtained from Amersham and is shown in Figure 14. An aliquot (2 µg) was digested with restriction enzymes SalI and EcoRI, the digest was run on a 1% agarose gel and the large DNA fragment (4520 bp) was purified. Two oligonucleotides were synthesized which, when annealed together, encode the OmpA polylinker region shown in Figure 15. This sequence has cohesive ends which are compatible with the SalI and EcoRI ends generated by restriction of pTTQ9. By cloning this oligonucleotide 'cassette' into the pTTQ9 vector, the SalI site is not regenerated, but the

EcoRI site is maintained. The cassette encodes the first 13 amino acids of the signal sequence of the *E. coli* outer-membrane protein Omp-A, preceded by the Shine Dalgarno ribosome-binding site of the OmpA gene. In addition restriction sites for enzymes XbaI, MunI, StyI and SphI are present. The MunI and StyI sites are within the coding region of the OmpA signal sequence and are intended as the 5' cloning sites for insertion of genes. The two oligonucleotides which make up this cassette were annealed together by mixing at a concentration of 5 pmoles/ $\mu$ l and heating in a waterbath to 95°C for 3 minutes, then slow cooling to room temperature. The annealed sequence was then ligated into the SalI / EcoRI cut pTTQ9. The resulting plasmid intermediate, termed pTQOmp, was verified by DNA sequencing.

(b) Fragment Preparation and Ligation

Plasmid pTTO-1 was constructed by ligating one DNA fragment from plasmid pACYC184 to two fragments generated from pTQOmp. Plasmid pACYC184 was obtained from New England Biolabs, and a restriction map is shown in Figure 16. An aliquot (2  $\mu$ g) was digested to completion with restriction enzyme StyI, then treated with Mung Bean Nuclease; this treatment creates blunt ends by cutting back 5' base overhangs. Following phenol extraction and ethanol precipitation, the DNA was restricted with enzyme PvuII, generating fragments of 2348, 1081, 412 and 403 bp. The 2348 bp fragment was purified after agarose gel electrophoresis. This fragment encodes the tetracycline resistance marker and the p15A origin of replication. The fragment was then treated with calf intestinal alkaline phosphatase to remove 5' terminal phosphates, thereby preventing the self-ligation of this molecule.

An aliquot (2  $\mu$ g) of plasmid pTQOmp was digested with enzymes SspI and EcoRI, and the 2350 bp fragment was purified from unwanted fragments of 2040 bp and 170 bp following agarose gel electrophoresis; this fragment encodes the transcriptional terminator region and the lacI<sup>q</sup> gene. Another aliquot (2  $\mu$ g) of pTQOmp was digested with EcoRI and XmnI, generating fragments of 2289, 1670, 350 and 250 bp. The 350 bp fragment, encoding the tac promoter, OmpA signal sequence and multicloning site, was gel purified.

The three fragments were then ligated, using approximately equimolar amounts of each fragment, to generate the plasmid pTTO-1. All cloning junctions were verified by DNA sequencing. The restriction map of this plasmid is shown in Figure 17. Plasmid pTTO-2 was then created by insertion of DNA encoding the human Ig light chain kappa constant domain. This was obtained as a SphI - EcoRI restriction fragment from plasmid

pHC132, and inserted into the corresponding sites in pTTO-1. Plasmid pTTO-2 is shown in Figure 18.

Insertion of humanized hTNF40 variable regions into pTTO-2

5        The variable light chain region hTNF40gL1 (SEQ ID NO:8) was obtained by PCR 'rescue' from the corresponding vector for mammalian cell expression pMR10.1. The OmpA leader sequence replaces the native Ig leader. The sequence of the PCR primers is shown below:

10    5' primer:

CGCGCGGCAATTGCAGTGGCCTTGGCTGGTTTCGCTACCGTAGCGCAAG  
CTGACATTCAAATGACCCAGAGCCC (SEQ ID NO:79)

3' primer:    TTCAACTGCTCATCAGATGG (SEQ ID NO:80)

15

Following PCR under standard conditions, the product was purified, digested with enzymes MunI and SphI then gel purified. The purified fragment was then inserted into the MunI / SphI sites of pTTO-2 to create the light chain intermediate pTTO(hTNF40L).

20        The variable heavy chain region of gh3hTNF40.4 was obtained in the same way from the vector pGamma-4. The sequence of the PCR primers is shown below:

5' primer:

GCTATCGCAATTGCAGTGGCGCTAGCTGGTTTCGCCACCGTGGCGCAAG  
CTGAGGTTTCAGCTGGTCGAGTCAGGAGGC (SEQ ID NO:81)

25

3' primer:    GCCTGAGTTCCACGACAC (SEQ ID NO:82)

30        Following PCR the product was purified, digested with enzymes NheI and ApaI then sub-cloned into the vector pDNAbEng-G1 (Figure 19). After verification by DNA sequencing, the heavy chain was restricted with enzyme EcoRI and sub-cloned into the EcoRI site of pTTO(hTNF40L) to create the *E. coli* expression plasmid pTTO(hTNF40).

### Optimisation of Intergenic Sequence for Modified Fab Expression

In the pTTO vector, modified Fab expression occurs from a dicistronic message encoding first light chain then heavy chain. The DNA sequence between the two genes (intergenic sequence, IGS) can influence the level of expression of the heavy chain by affecting the rate of translational initiation. For example, a short intergenic sequence may result in translational coupling between the light and heavy chains, in that the translating ribosome may not fully dissociate from the mRNA after completing light chain synthesis before initiating heavy chain synthesis. The 'strength' of any Shine Dalgarno (SD) ribosome binding site (homology to 16S rRNA) can also have an effect, as can the distance and sequence composition between the SD and the ATG start codon. The potential secondary structure of mRNA around the ATG is another important factor; the ATG should be in a 'loop' and not constrained within a 'stem', while the reverse applies to the SD. Thus by modifying the composition and length of the IGS it is possible to modify the strength of translational initiation and therefore the level of heavy chain production. It is likely that an optimum rate of translational initiation needs to be achieved to maximise expression of the heavy chain of a given modified Fab. For example, with one modified Fab, a high level of expression may be tolerated, but for a different modified Fab with different amino acid sequence, a high level of expression might prove toxic, perhaps because of different efficiencies of secretion or folding. For this reason, a series of four intergenic sequences were designed (Figure 20), permitting the empirical determination of the optimum IGS for the hTNF40-based modified Fab. IGS1 and IGS2 have very short intergenic sequences (-1 and +1 respectively) and might be expected to give closely coupled translation; the SD sequences (underlined) are subtly different. These two sequences will most likely confer a high level of translational initiation. IGS3 and IGS4 have a longer distance between start and stop codons (+13) and differ in their sequence composition; IGS3 has a 'stronger' SD sequence. All sequences were studied for secondary structure (using m/fold program) and 'optimised' as far as possible; however, with tight coupling of translation of the two chains the lack of ribosomal dissociation means that the mRNA may not be 'naked', preventing secondary structure formation.

### Cloning of IGS variants

The IGS cassettes shown in Figure 20 have flanking SacI and MunI cloning sites. They were built by annealing complementary oligonucleotide pairs. A vector fragment was prepared by digesting pTTO(hTNF40) with SacI and NotI, and a heavy chain fragment was



prepared by digesting pDNAbEngG1(hTNF40H) with MunI and NotI. Three-way ligations were then performed, using equimolar amounts of the two restriction fragments and approximately 0.05 pmoles of each annealed oligo cassette. This created the four expression plasmids pTTO(hTNF40 IGS-1), pTTO(hTNF40 IGS-2), pTTO(hTNF40 IGS-3), pTTO(hTNF40 IGS-4).

#### **Shake flask expression analysis**

The four plasmids were transformed into *E. coli* strain W3110, along with the original expression construct, and then analysed for expression in shake flasks as described.

10 The results of a typical experiment are shown in Figure 21. The different intergenic sequences confer different expression profiles. IGS1 and IGS2 accumulate periplasmic modified Fab rapidly with a peak at 1 hour post induction, after which the level recovered falls. The peak is greater and the fall sharper for IGS1. These results are consistent with a high level of synthesis, as expected for close translational coupling for these constructs.

15 IGS1 apparently confers a higher level of heavy chain expression than does IGS2. In this instance, it appears that this high level of expression is poorly tolerated, since periplasmic expression levels fall after the 1 hour peak. This is seen on the growth profile of the IGS1 culture (not shown), which peaks at 1 hour post induction before falling, suggesting cell death and lysis. IGS3 accumulates modified Fab more slowly but peaks at 2 hours post

20 induction with a higher peak value (325 ng/ml/OD), before levels fall. The growth of this culture continued to 3 hours post induction and reached a higher peak biomass (not shown). This is consistent with a lower level of heavy chain synthesis. IGS4 accumulates material at a slower rate still and fails to reach the high peak of productivity of the other 3 constructs. All IGS variants out-perform the original vector significantly. The hypothesis

25 that the different IGS sequences confer different rates of translational initiation is supported by these experimental results. For the hTNF40-based modified Fab it appears that a high rate of heavy chain translational initiation is poorly tolerated and is therefore not optimal. A slower rate, as conferred by IGS3, results in better growth characteristics and consequently a better yield accumulates over time.

30 Following comparison of productivity in the fermenter the IGS3 construct was selected as the highest yielding and was termed pTTO(CDP870) – see Figure 22.

The heavy chain encoded by the plasmid pTTO(CDP870) has the sequence given in SEQ ID NO:115 and the light chain has the sequence given in SEQ ID NO:113.

### PEGylation of CDR-Grafted, hTNF40-based Modified Fab.

The purified modified Fab is site-specifically conjugated with a branched molecule of PEG. This is achieved by activation of a single cysteine residue in a truncated hinge region of the modified Fab, followed by reaction with (PEG)-lysyl maleimide as previously described (A.P. Chapman *et al.*, Nature Biotechnology 17, 780-783; 1999). The PEGylated molecule is shown in Figure 13 and is called compound CDP870:

### Efficacy of PEGylated CDR-Grafted, hTNF40-based Modified Fab (CDP870) in Treating Rheumatoid Arthritis.

CDP870 has a long half life of approximately 11 days.

We evaluated the safety and efficacy of intravenous CDP870 in a randomised double-blind placebo-controlled dose escalating trial in patients with RA.

### Methods

#### 15 Patients:

Patients aged between 18 and 75 years old and who satisfied the 1987 revised American College of Rheumatology (ACR) diagnostic criteria for rheumatoid arthritis (RA) (Arnett *et al.*, Arthritis Rheum., 31, 315-324, 1988) were recruited from outpatient Rheumatology clinics at London, Cambridge, Norfolk and Norwich (United Kingdom). Patients were required to have clinically active disease as defined by having at least 3 of the following criteria:  $\geq 6$  painful or tender joints;  $\geq 45$  minutes of early morning stiffness; and erythrocyte sedimentation rate (ESR)  $\geq 28$  mm/hr. They must have failed to respond to at least one Disease Modifying Anti-Rheumatic Drug (DMARD) and have been off treatment for at least 4 weeks. Corticosteroids were permitted if the dose was  $\geq 7.5$  mg/day of prednisolone. Pregnant women, nursing women and women of childbearing potential not using an effective method of contraception were excluded. Patients were also excluded if they had a previous history of malignancy, concomitant severe uncontrolled medical conditions, previous failure of TNF $\alpha$ -neutralizing therapy or allergy to polyethylene glycol. Written informed consent was obtained from each patient before enrolment. The study was approved by the local research ethics committees.

*Treatment protocol:*

36 RA patients were divided into 3 groups, each to receive an increasing dose of the trial drug (1, 5 or 20mg/kg). Each group of 12 was randomly divided into 8 to receive CDP870 and 4 to receive placebo. CDP870 was given as a single intravenous infusion (100 ml in total) over 60 minutes. Placebo (sodium acetate buffer) was given similarly as a single intravenous infusion of 100 ml over 60 minutes. Treatment was given on an outpatient basis. After 8 weeks, all patients had the opportunity to receive an infusion of either 5 or 20 mg/kg of CDP870 in open fashion.

10 *Clinical assessment:*

RA disease activity was assessed based on the World Health Organization and International League of Associations for Rheumatology (Boers *et al.*, J. Rheumatol – Supplement, 41, 86-89, 1994) and European League Against Rheumatism (EULAR) (Scott *et al.*, Clin. Exp. Rheumatol., 10, 521-525, 1992) core data sets with 28 joint counts. Changes in disease activity were assessed by Disease Activity Score (Prevoo *et al.*, Arthritis Rheum., 38, 44-48, 1995) and the ACR responses criteria (Felson *et al.*, Arthritis Rheum., 38, 727-735, 1995). Assessments were carried out before treatment and at 1, 2, 4, 6 and 8 weeks after therapy. Patients were also assessed for safety and tolerance of the study drug. Haematology, biochemistry, anti-CDP870 antibodies and adverse events were assessed at each visit.

*CDP870 plasma concentration and anti-CDP870 antibodies:*

CDP870 was measured by enzyme-linked immunosorbent assay (ELISA). Serial dilutions of patients' plasma were incubated in microtitre plates (Nunc) coated with recombinant human TNF $\alpha$  (Strathmann Biotech GmbH, Hannover). Captured CDP870 was revealed with horseradish peroxidase conjugated goat anti-human kappa light chain (Cappel, ICN) followed by tetramethylbenzidine (TMB) substrate.

Antibodies to CDP870 was screened (at 1/10 plasma dilution) using a double antigen sandwich ELISA with biotinylated CDP870 as the second layer. Bound antibodies were revealed using HRP-streptavidin and TMB substrate. The assay was calibrated using a hyperimmune rabbit IgG standard. A unit of activity is equivalent to 1  $\mu$ g of the rabbit standard.

### Statistical Analysis

The study was exploratory in nature and the sample size was based on previous experience with similar agents. Efficacy of CDP870 was analysed by calculating disease activity score (DAS) and ACR20/50 responses for intention to treat and per-protocol using a closed testing procedure. The disease activity score was calculated as follows:  $DAS = 0.555 \times \text{square root of (28 tender joints)} + 0.284 \times \text{square root of (28 swollen joints)} + 0.7 \times \ln(\text{ESR}) + 0.0142 \times (\text{patient's global assessment})$ . First, the pooled active groups were compared to placebo. If this comparison was significant at the 5% level, each dosage group was compared to placebo. All comparisons were two tailed with a significance level of 5%. All P-values were derived from exploratory analysis and should not be used for inferential interpretation.

### Results

#### Demography:

36 patients with RA were recruited. Their demographic details are given in Table 6. The mean age was 56 years and 30 patients were female. The mean duration of RA was 13 years and 21 patients were rheumatoid factor positive. Patients in the different groups have similar demographic characteristics. In the blinded dosing period, 6/12 placebo-treated patients withdrew from the study for deteriorating RA  $\geq 4$ , weeks after dosing. 2/24 CDP870-treated patients withdrew, both in the 1 mg/kg group, for deteriorating RA/lost to follow up  $> 4$  weeks after dosing. The difference was statistically significant ( $p=0.009$ , Fisher exact test).

**Table 6: Demographic details (mean  $\pm$  standard deviation)**

	Number	Sex (M:F)	Age	Duration of Disease	Rheuma- toid Factor	Number of previous DMARDs
Placebo	12	1:11	51 $\pm$ 8	12 $\pm$ 8	8(67%)	5 $\pm$ 1
1 mg/kg	8	1:7	59 $\pm$ 7	12 $\pm$ 7	4(50%)	4 $\pm$ 1
5m g/kg	8	2:6	54 $\pm$ 13	13 $\pm$ 5	5(63%)	5 $\pm$ 2
20 mg/kg	8	2:6	61 $\pm$ 11	14 $\pm$ 13	4(50%)	4 $\pm$ 2

*Clinical Efficacy:*

The proportion of patients with ACR20 improvement for the per-protocol population with last observation carried forward was 16.7, 50, 87.5 and 62.5% after placebo, 1, 5 and 20 mg/kg CDP870 (combined treatment effect  $p=0.012$ ) at 4 weeks and 16.7, 25, 75 and 75% ( $p=0.032$ ) at 8 weeks. Reduction in DAS scores (median) for the per-protocol population with last observation carried forward was 0.15, 1.14, 1.91 and 1.95 after placebo, 1, 5 and 20 mg/kg CDP870 (combined treatment effect  $p=0.001$ ) at 4 weeks and 0.31, 0.09, 2.09 and 1.76 ( $p=0.008$ ) at 8 weeks (Figure 23). Changes in individual components of the World Health Organization and International League of Associations for Rheumatology core data set are shown in Figure 24.

Following the open label dose of CDP870, similar beneficial effects were achieved. Of the 36 patients recruited into the study, 32 received a second infusion of CDP870. The proportion of patients with ACR20 improvement from pre-first infusion was 72.2 and 55.6% after 5 and 20 mg/kg CDP870 at 4 weeks and 55.6 and 66.7% at 8 weeks.

*Adverse Events*

Treatment was well tolerated with no infusion-related reaction. No allergic reaction or skin rash was reported. In the double-blind phase, there were 19, 38, 8 and 14 adverse events in the placebo, 1, 5 and 20 mg/kg groups respectively. The commonest was headache with 9 episodes in 5 patients (1 placebo, 3 at 1 mg/kg, 1 at 20 mg/kg). One patient who received placebo and 3 patients who received CDP870 (1 at 5 mg/kg and 2 at 20 mg/kg) developed lower respiratory tract infections. These were reported as mild or moderate. They were treated with oral antibiotics and resolved over 1-2 week period. Three patients each in the 1 and 5 mg/kg groups and one in the 20 mg/kg group developed a urinary tract infection 1-2 months after CDP870 treatment. One adverse event was described as severe which was an episode of neck pain occurring 3 days after infusion with 1 mg/kg. Increase in anti-nuclear antibody was seen in 4 patients: 1 in the placebo group (negative to 1/40), 2 in the 1 mg/kg group (negative to 1/40, negative to 1/80) and 1 in the 20 mg/kg group (negative to 1/40). No change was found in anti-DNA or anti-cardiolipin antibodies.

### CDP870 Plasma Concentration and Anti-CDP870 levels

As expected, for all dose levels of CDP870, the peak plasma concentration occurred at the end of infusion and was dose proportional with plasma concentration declining slowly thereafter. The plasma concentration profile of CDP870 appeared very similar to that previously observed in volunteers where the half-life was calculated to be approximately 14 days. On re-dosing, a similar profile to single dose infusion was observed.

Following a single intravenous infusion, anti-CDP870 levels were low or undetectable.

### Discussion

Neutralizing TNF $\alpha$  is an effective treatment strategy in RA. Currently, this requires the use of biological agents, such as a chimeric mAb or a soluble receptor/human Fc fusion protein, which are expensive to manufacture. A therapeutic TNF $\alpha$  neutralizing agent needs to bind TNF $\alpha$  with high affinity and have a long plasma half-life, low antigenicity and high tolerability and safety. It also needs to be accessible to all patients with RA who would benefit from TNF $\alpha$  blockade. One technology that could achieve these objectives is the conjugation with polyethelene glycol of a TNF $\alpha$  binding antibody fragment made in *E. coli*. In this preliminary study, we find that CDP870, a PEGylated, anti-TNF $\alpha$ , modified Fab, is effective and well tolerated by patients with RA.

*In vitro* studies have shown that CDP870 has similar TNF $\alpha$  neutralizing activity to the murine anti-TNF $\alpha$  parent antibody. This study confirms that CDP870 reduced inflammation and improved symptoms in RA. Clinical improvement as measured by the ACR20 response criteria in the 5 and 20 mg/kg groups (75%, 75%) was comparable to etanercept (60%) (Moreland *et al.*, *Annals Int. Med.*, 130, 478-486, 1999) and infliximab (50%) (Maini *et al.*, *Lancet*, 354, 1932-1939, 1999). At the middle and highest dosage levels tested, the therapeutic effect lasted 8 weeks which is comparable to previous other mAbs (Elliott *et al.*, *Lancet*, 344, 1105-1110, 1994 and Rankin *et al.*, *Br. J. Rheumatol.*, 34, 334-342, 1995). Previous study has shown that the therapeutic effect of anti-TNF $\alpha$  antibody is related to its plasma half-life and the generation of circulating antibodies (Maini *et al.*, *Arthritis Rheum.* 38 (Supplement) : S186 1995 (Abstract)). Our study showed that CDP870 has a plasma half-life of 14 days which is equivalent to that of a whole antibody

(Rankin *et al.*, (*supra*)) and much longer than the half-life of unconjugated Fab' fragments. Further, CDP870 generated only very low levels of antibody response.

One of the important objectives of this study is to examine the tolerability and safety of administering this PEGylated Fab'. In our study, CDP870 appears well tolerated.  
5 Although further study will be needed to assess long-term toxicity, especially the risk of demyelinating disease, infection and skin rashes that have been reported with etanercept and infliximab.

In summary, CDP870 is therapeutically effective in RA and was well tolerated in this short-term study.

10 It should be understood that the above-described examples are merely exemplary and do not limit the scope of the present invention as defined in the following claims.

## CLAIMS

1. An antibody molecule having specificity for human TNF $\alpha$ , comprising a heavy  
5 chain wherein the variable domain comprises a CDR having the sequence given as H1 in  
Figure 3 (SEQ ID NO:1) for CDRH1, as H2' in Figure 3 (SEQ ID NO:2) or as H2' in  
Figure 3 (SEQ ID NO:7) for CDRH2 or as H3 in Figure 3 (SEQ ID NO:3) for CDRH3.
2. An antibody molecule having specificity for human TNF $\alpha$ , comprising a light  
10 chain wherein the variable domain comprises a CDR having the sequence given as L1 in  
Figure 3 (SEQ ID NO:4) for CDRL1, as L2 in Figure 3 (SEQ ID NO:5) for CDRL2 or as  
L3 in Figure 3 (SEQ ID NO:6) for CDRL3.
3. The antibody molecule of claim 1 or claim 2 comprising a heavy chain wherein the  
15 variable domain comprises a CDR having the sequence given in SEQ ID NO:1 for CDRH1,  
SEQ ID NO:2 or SEQ ID NO:7, for CDRH2 or SEQ ID NO:3 for CDRH3 and a light chain  
wherein the variable domain comprises a CDR having the sequence given in SEQ ID NO:4  
for CDRL1, SEQ ID NO:5 for CDRL2 or SEQ ID NO:6 for CDRL3.
- 20 4. The antibody molecule of claim 3, which comprises SEQ ID NO:1 for CDRH1,  
SEQ ID NO: 2 or SEQ ID NO:7 for CDRH2, SEQ ID NO:3 for CDRH3, SEQ ID NO:4 for  
CDRL1, SEQ ID NO:5 for CDRL2 and SEQ ID NO:6 for CDRL3.
5. The antibody molecule of any one of claims 1 to 4, which comprises SEQ ID NO:2  
25 for CDRH2.
6. The antibody molecule of any one of claims 1 to 5, which is a CDR-grafted  
antibody molecule.
- 30 7. The antibody molecule of claim 6, wherein the variable domain comprises human  
acceptor framework regions and non-human donor CDRs.



8. The antibody molecule of claim 7, wherein the human acceptor framework regions of the variable domain of the heavy chain are based on a human group 1 consensus sequence and comprise non-human donor residues at positions 28, 69 and 71
- 5 9. The antibody molecule of claim 7, wherein the human acceptor framework regions of the variable domain of the heavy chain are based on a human group 1 consensus sequence and comprise non-human donor residues at positions 28, 38, 46, 67, 69 and 71.
- 10 10. The antibody molecule of claim 7, wherein the human acceptor framework regions of the variable domain of the heavy chain are based on a human group 3 consensus sequence and comprise non-human donor residues at positions 27, 28, 30, 48 49, 69, 71, 73 76 and 78.
- 15 11. The antibody molecule of any one of claims 7 to 10, wherein the human acceptor framework regions of the variable domain of the light chain are based on human group 1 consensus sequence and comprise non-human donor residues at positions 46 and 60.
- 20 12. The antibody molecule of any one of claims 1 to 11, comprising the light chain variable region hTNF40-gL1 (SEQ ID NO:8) and the heavy chain variable region gh3hTNF40.4 (SEQ ID NO:11).
13. The antibody molecule of any one of claims 1 to 12 which is a Fab fragment.
- 25 14. The antibody molecule of claims 12 and 13, which is a Fab fragment comprising a heavy chain having the sequence given in SEQ ID NO:111 and a light chain having the sequence given in SEQ ID NO:113.
- 30 15. The antibody molecule of any one of claims 1 to 12, which is a modified Fab fragment having at the C-terminal end of its heavy chain one or more amino acids to allow attachment of an effector or reporter molecule.

16. The antibody molecule of claim 15, wherein the additional amino acids form a modified hinge region containing one or two cysteine residues to which the effector or reporter molecule may be attached.
- 5 17. The antibody molecule of claim 12, which is a modified Fab fragment comprising a heavy chain having the sequence given in SEQ ID NO:115 and a light chain having the sequence given in SEQ ID NO:113.
- 10 18. An antibody molecule having specificity for human TNF $\alpha$ , having a light chain comprising the sequence given in SEQ ID NO:113.
19. An antibody molecule having specificity for human TNF $\alpha$ , having a light chain consisting of the sequence given in SEQ ID NO:113.
- 15 20. An antibody molecule having specificity for human TNF $\alpha$ , having a heavy chain comprising the sequence given in SEQ ID NO:115.
21. An antibody molecule having specificity for human TNF $\alpha$ , having a heavy chain consisting of the sequence given in SEQ ID NO:115.
- 20 22. An antibody molecule having specificity for human TNF $\alpha$ , having a light chain comprising the sequence given in SEQ ID NO:113 and a heavy chain comprising the sequence given in SEQ ID NO:115.
- 25 23. An antibody molecule having specificity for human TNF $\alpha$ , having a light chain consisting of the sequence given in SEQ ID NO:113 and a heavy chain consisting of the sequence given in SEQ ID NO:115.
24. A variant of the antibody molecule of any one of claims 1 to 23, which has an improved affinity for TNF $\alpha$ .
- 30 25. The variant of claim 24 which is obtained by an affinity maturation protocol.

26. The antibody of any one of claims 1 to 3 which is murine anti-TNF $\alpha$  monoclonal antibody hTNF40.
27. The antibody molecule of any one of claims 1 to 3, which is a chimeric antibody  
5 molecule comprising the light and heavy chain variable domains of the monoclonal antibody of claim 26.
28. A compound comprising the antibody molecule of any one of claims 15 to 23 having covalently attached to an amino acid at or towards the C-terminal end of its heavy  
10 chain an effector or reporter molecule.
29. The compound of claim 28, which comprises an effector molecule.
30. The compound of claim 29, wherein the effector molecule comprises one or more  
15 polymers.
31. The compound of claim 30, wherein the one or more polymers is/are an optionally substituted straight or branched chain polyalkylene, polyalkenylene or polyoxyalkylene polymer or a branched or unbranched polysaccharide.  
20
32. The compound of claim 31, wherein the one or more polymers is/are a methoxypoly(ethyleneglycol).
33. A compound comprising the antibody molecule of claim 17 having attached to one  
25 of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide group wherein each amino group of the lysyl residue has covalently linked to it a methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da.
34. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
30 having a light chain comprising the sequence given in SEQ ID NO:113 and a heavy chain comprising the sequence given in SEQ ID NO:115, having attached to one of the cysteine residues at the C-terminal end of the heavy chain one or more synthetic or naturally-occurring polymers.

35. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
having a light chain consisting of the sequence given in SEQ ID NO:113 and a heavy chain  
consisting of the sequence given in SEQ ID NO:115, having attached to one of the cysteine  
5 residues at the C-terminal end of the heavy chain one or more synthetic or naturally-  
occurring polymers.
36. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
having a light chain comprising the sequence given in SEQ ID NO:113, having attached to  
10 one of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide  
group wherein each amino group of the lysyl residue has covalently linked to it a  
methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da..
37. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
15 having a light chain consisting of the sequence given in SEQ ID NO:113, having attached  
to one of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide  
group wherein each amino group of the lysyl residue has covalently linked to it a  
methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da.
- 20 38. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
having a heavy chain comprising the sequence given in SEQ ID NO:115, having attached  
to one of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide  
group wherein each amino group of the lysyl residue has covalently linked to it a  
methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da.  
25
39. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
having a heavy chain consisting of the sequence given in SEQ ID NO:115, having attached  
to one of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide  
group wherein each amino group of the lysyl residue has covalently linked to it a methoxy  
30 poly(ethyleneglycol) residue having a molecular weight of about 20,000 Da.
40. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ ,  
having a light chain comprising the sequence given in SEQ ID NO:113 and a heavy chain

comprising the sequence given in SEQ ID NO:115, having attached to one of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide group wherein each amino group of the lysyl residue has covalently linked to it a methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da.

5

41. A compound comprising an antibody molecule having specificity for human TNF $\alpha$ , having a light chain consisting of the sequence given in SEQ ID NO:113 and a heavy chain consisting of the sequence given in SEQ ID NO:115, having attached to one of the cysteine residues at the C-terminal end of the heavy chain a lysyl-maleimide group wherein each  
10 amino group of the lysyl residue has covalently linked to it a methoxypoly(ethyleneglycol) residue having a molecular weight of about 20,000 Da.

42. An antibody molecule comprising a hybrid CDR comprising a truncated donor CDR sequence wherein the missing portion of the donor CDR is replaced by a different sequence  
15 and forms a functional CDR.

43. The antibody molecule of claim 42, wherein the missing part of the CDR sequence is from the antibody from which the framework regions of the antibody molecule are derived.

20

44. The antibody molecule of claim 43, wherein the missing part of the CDR sequence is derived from a germline antibody having consensus framework regions.

45. The antibody molecule of any one of claims 42 to 44, wherein CDRH2 of the heavy  
25 chain is hybrid in the antibody molecule.

46. The antibody molecule of any one of claims 42 to 45, wherein the truncation of the donor CDR is from 1 to 8 amino acids.

30 47. The antibody molecule of claim 46, wherein the truncation is from 4 to 6 amino acids.

48. The antibody molecule of any one of claims 42 to 47, wherein the truncation is made at the C-terminus of the CDR.
- 
49. A DNA sequence which encodes the heavy and/or light chain of the antibody molecule of any one of claims 1 to 27 and 42 to 48.
50. The DNA sequence of claim 49 comprising the sequence shown in SEQ ID NO:8 or 10.
51. The DNA sequence of claim 49 comprising the sequence shown in SEQ ID NO:10 or 11.
52. The DNA sequence of claim 49 comprising the sequence shown in SEQ ID NO:110, 112 or 114.
53. A cloning or expression vector containing the DNA sequence of any one of claims 49 to 52.
54. An *E. coli* expression vector comprising the DNA sequence of any one of claims 49 to 52.
55. The *E. coli* expression vector of claim 54 which is pTTO(CDP870).
56. A host cell transformed with the vector of any one of claims 53 to 55.
57. A process for the production of the antibody molecule of any one of claims 1 to 27 and 42 to 48, comprising culturing the host cell of claim 56 and isolating the antibody molecule.
58. A process for the production of the antibody molecule of any one of claims 1 to 27 and 42 to 48, comprising culturing *E. coli* comprising an *E. coli* expression vector comprising the DNA sequence of any one of claims 53 to 55 and isolating the antibody molecule.

59. The process of claim 58 wherein the antibody molecule is targeted to the periplasm.
60. A therapeutic or diagnostic composition comprising the antibody molecule of any  
5 one of claims 1 to 27 and 42 to 48 or the compound of any one of claims 28 to 41.
61. The antibody molecule of any one of claims 1 to 27 and 42 to 48, having specificity  
for human TNF $\alpha$ , or the compound of any one of claims 28 to 41, for use in treating a  
pathology mediated by TNF $\alpha$ .
- 10 62. The antibody molecule of or compound claim 61, for use in treating rheumatoid- or  
osteo- arthritis.
63. Use of the antibody molecule of any one of claims 1 to 27 and 42 to 48, having  
15 specificity for human TNF $\alpha$ , or the compound of any one of claims 28 to 41 in the  
manufacture of a medicament for the treatment of a pathology mediated by TNF $\alpha$ .
64. The use of claim 63, wherein the pathology is rheumatoid- or osteo- arthritis.
- 20 65. The vector pDNA<sub>AbEng-G1</sub> as shown in Figure 19.
66. The vector pTTO(CDP870) as shown in Figure 22.
67. A polypeptide having the amino acid sequence given in any one of SEQ ID NOS:1  
25 to 7.

1/27

**FIG. 1****Comparisons of framework regions of light chain of antibody hTNF40 and human group 1 consensus sequences**

Hu group 1 consensus : DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO: 83)

hTNF40 : DIVMTQSOKFMSTSVGDRVSVTC (SEQ ID NO: 84)

Hu Group 1 consensus : WYQOKPGKAPKLLIY (SEQ ID NO: 85)

hTNF40 : WYQOKPGQSPKALIY (SEQ ID NO: 86)Hu Group 1 consensus : GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO: 87)hTNF40 : GVPYRFTGSGSGTDFTLTISTVQSEDLAEYFC (SEQ ID NO: 88)

Hu Group 1 consensus : FGQGTKVEIKR (SEQ ID NO: 89)

hTNF40 : FGAGTKLELKR (SEQ ID NO: 90)**FIG. 3 Sequence of CDRs of hTNF40**

H1 DYGMN (SEQ ID NO:1)

H2 WINTYIGEPIYVDDFKG (SEQ ID NO:7)

H2' WINTYIGEPIYADSVKG (SEQ ID NO:2)

H3 GYRSYAMDY (SEQ ID NO:3)

L1 KASQNVGTNVA (SEQ ID NO:4)

L2 SASFLYS (SEQ ID NO:5)

L3 QQYNIYPLT (SEQ ID NO:6)



2/27

## FIG. 2

Comparisons of framework regions of heavy chain of antibody hTNF40 and human group 1 and group 3 consensus sequences

Hu Group 1 consensus : QVQLVQSGAEVKKPGASVKVSCKASGYTFT (SEQ ID NO: 91)  
hTNF40 : QIQLVQSGPELKKPGETVKISKASGYVFT (SEQ ID NO: 92)

Hu Group 1 consensus : WVRQAPGQGLEWMG (SEQ ID NO: 93)  
hTNF40 : WVKQAPGKAFKWMG (SEQ ID NO: 94)

Hu Group 1 consensus : RVTITRDTSTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO: 95)  
hTNF40 : RFAFSLETSASTAFLQINNLKNEDTATYFCAR (SEQ ID NO: 96)

Hu Group 1 consensus : WGQGTLLTVSS (SEQ ID NO: 97)  
hTNF40 : WGQGTLLTVSS (SEQ ID NO: 98)

Hu Group 3 consensus : EVQLVESGGGLVQPGGSLRLSCAASGFTFS (SEQ ID NO: 106)  
hTNF40 : QIQLVQSGPELKKPGETVKISKASGYVFT (SEQ ID NO: 92)

Hu Group 3 consensus : WVRQAPGKGLEWVS (SEQ ID NO: 107)  
hTNF40 : WVKQAPGKAFKWMG (SEQ ID NO: 94)

Hu Group 3 consensus : RFTISRDNKNTLYLQMNSLRAEDTAVYYCAR (SEQ ID NO: 108)  
hTNF40 : RFAFSLETSASTAFLQINNLKNEDTATYFCAR (SEQ ID NO: 96)

Hu Group 3 consensus : WGQGTLLTVSS (SEQ ID NO: 109)  
hTNF40 : WGQGTLLTVSS (SEQ ID NO: 98)

3/27  
FIG. 4





FIG. 5



4/27

FIG. 6 Murine VI Sequence of hTNF40 (SEQ ID NO: 99)

```

10      20      30      40      50
GAC ATT GTG ATG ACC CAG TCT CAA AAA TTC ATG TCC ACA TCA GTA GGA GAC AGG
CTG TAA CAC TAC TGG GTC AGA GTT TTT AAG TAC AGG TGT AGT CAT CCT CTG TCC
D   I   V   M   T   Q   S   Q   K   F   M   S   T   S   V   G   D   R>

60      70      80      90      100
GTC AGC GTC ACC TGC AAG GCC AGT CAG AAT GTG GGT ACT AAT GTA GCC TGG TAT
CAG TCG CAG TGG ACG TTC CGG TCA GTC TTA CAC CCA TGA TTA CAT CGG ACC ATA
V   S   V   T   C   K   A   S   Q   N   V   G   T   N   V   A   W   Y>

110     120     130     140     150     160
CAA CAG AAA CCA GGA CAA TCT CCT CCA AAA GCA CTG ATT TAC TCG GCA TCC TTC CTA
GTT GTC TTT GGT CCT GGT AGA GGA TTT CGT GAC TAA ATG AGC CGT AGG AAG GAT
Q   Q   K   P   G   Q   S   S   P   K   A   L   I   Y   S   A   S   F   L>

170     180     190     200     210
TAT AGT GGA GTC CCT TAT CGC TTC ACA GGC AGT GGA TCT GGG ACA GAT TTC ACT
ATA TCA CCT CAG GGA ATA GCG AAG TGT CCG TCA CCT AGA CCC TGT CTA AAG TGA
Y   S   G   V   P   Y   R   F   T   G   S   G   S   G   T   D   F   T>

220     230     240     250     260     270
CTC ACC ATC AGC ACT GTG CAG TCT GAA GAC TTG GCA GAG TAT TTC TGT CAG CAA
GAG TGG TAG TCG TGA CAC GTC AGA CTT CTG AAC CGT CTC ATA AAG ACA GTC GTT
L   T   I   S   T   V   Q   S   E   D   L   A   E   Y   F   C   Q   Q>

280     290     300     310     320
TAT AAC ATC TAT CCT CTC ACG TTC GGT GCT GGG ACC AAG CTG GAG CTG AAA CGT
ATA TTG TAG ATA GGA GAG TGC AAG CCA CGA CCC TGG TTC GAC CTC GAC TTT GCA
Y   N   I   I   Y   P   L   T   F   G   A   G   T   K   L   E   L   K   R>

```

5/27

FIG. 7 Murine Vh Sequence of hTNF40 (SEQ ID NO: 100)

```

10  CAG ATC CAG TTG GTG CAG TCT GGA CCT GAG AAG AAG CCT GGA GAG ACA GTC
20  GTC TAG GTC AAC CAC CAC GTC AGA CCT GGA CTC GAC TTC TTC GGA CCT CTC TGT CAG
30  Q I Q L V Q S G P E L K K P G E T V>
40
50
60  AAG ATC TCC TGC AAG GCT TCT GGA TAT GTT TTC ACA GAC TAT GGA ATG AAT TGG
70  TTC TAG AAG ACG TTC CGA AGA CCT ATA CAA AAG TGT CTG ATA CCT TAC TTA ACC
80  K I S C K A S G Y V F T D Y G M N W>
90
100
110 GTG AAG CAG GCT CCA GGA AAG GCT TTC AAG TGG ATG GGC TGG ATA AAC ACC TAC
120 CAC TTC GTC CGA GGT CCT TTC CGA AAG TTC ACC TAC CCG ACC TAT TAT TGG TGG ATG
130 V K Q A P G K A F K W M G W I N T Y>
140
150
160
170 ATT GGA GAG CCA ATA TAT GTT GAT GAC TTC AAG GGA CGA TTT GCC TTC TCT TTG
180 TAA CCT CTC GGT TAT ATA CAA CTA CTG AAG TTC CCT GCT AAA CGG AAG AGA AAC
190 I G E P I Y V D D F K G R F A F S L>
200
210
220 GAA ACC TCT GCC AGC ACT GCC TTT TTG CAG ATC AAC AAC CTC AAA AAT GAG GAC
230 CTT TGG AGA CGG TCG TGA CGG AAA AAC GTC TAG TTG TTG GAG TTT TTA CTC CTG
240 E T S A S T A F L Q I N N L K N E D>
250
260
270
280 ACG GCT ACA TAT TTC TGT GCA AGA GGT TAC CGG TCC TAT GCT ATG GAC TAC TGG
290 TGC CGA TGT ATA AAG ACA CGT TCT CCA ATG GCC AGG ATA CGA TAC CTG ATG ACC
300 T A T Y F C A R G Y R S Y A M D Y W>
310
320
330 GGT CAA GGA ACC TCA GTC ACC GTC TCT TCA
340 CCA GTT CCT TGG AGT CAG TGG CAG AGA AGT
350 G Q G T S V T V S S>

```

6/27

FIG. 8 Grafted VI Sequence of hTNF40 (SEQ ID NO: 8)

```

10  GAC ATT CAA ATG ACC CAG AGC CCA TCC AGC CTG AGC GCA TCT GTA GGA GAC CGG
20  CTG TAA GTT TAC TGG GTC TCG AGG TCG GAC TCG CGT AGA CAT CCT CTG GCC
30  D I Q M T Q S P S S L S A S V G D R>
40
50
60  GTC ACC ATC ACT TGT AAA GCC AGT CAG AAC GTA GGT ACT AAC GTA GCC TGG TAT
70  CAG TGG TAG TGA ACA TTT CGG TCA GTC TCG CAT CCA TGA TTG CAT CGG ACC ATA
80  V T I T C K A S Q N V G T N V A W Y>
90
100
110  CAG CAA AAA CCA GGT AAA GCC CCA AAA GCC CTC ATC TAC AGT GCC TCT TTC CTC
120  GTC GTT TTT GGT CCA TTT CGG GGT TTT CGG GAG TAG ATG TCA CGG AGA AAG GAG
130  Q Q K P G K A P K A L I Y S A S F L>
140
150
160
170  TAT AGT GGT GTA CCA TAC TAC AGG TTC AGC GGA TCC GGT AGT GGT ACT GAT TTC ACC
180  ATA TCA CCA CAT GGT ATG TCC AAG TCG CCT AGG CCA TCA CCA TGA CTA AAG TGG
190  Y S G V P Y R F S G S G T D F T>
200
210
220  CTC ACG ATC AGT AGC CTC CAG CCA GAA GAT TTC GCC ACT TAT TAC TGT CAA CAG
230  GAG TGC TAG TCA TCG GAG GTC GGT CTT CTA AAG CGG TGA ATA ATG ACA GTT GTC
240  L T I S S L Q P E D F A T Y Y C Q Q>
250
260
270
280  TAT AAC ATC TAC CCA CTC ACA TTC GGT CAG GGT ACT AAA GTA GAA ATC AAA
290  ATA TTG TAG ATG GGT GAG TGT AAG CCA GTC CCA TGA TTT CAT CTT TAG TTT
300  Y N I Y P L T F G Q G T K V E I K>
310
320

```

7/27

FIG. 9 Grafted VI sequence of hTNF40 (SEQ ID NO: 9)

```

10      GAC ATT CAA ATG ACC CAG AGC CCA TCC AGC CTG AGC GCA TCT GTA GGA GAC CGG
      CTG TAA GTT TAC TGG GTC TCG GGT AGG TCG GAC TCG CGT AGA CAT CCT CTG GCC
      D   I   Q   M   T   Q   S   P   S   S   L   S   A   S   V   G   D   R>

      20      30      40      50

60      GTC ACC ATC ACT TGT AAA GCC AGT CAG AAC GTA GGT ACT AAC GTA GCC TGG TAT
      CAG TGG TAG TGA ACA TTT CGG TCA GTC TTT CAT CCA TGA TTG CAT CGG ACC ATA
      V   T   I   T   C   K   A   S   Q   N   V   G   T   N   V   A   W   Y>

      70      80      90      100

110     GTC CAA AAA CCA CCA GGT AAA GCC CCA AAA CTC CTC ATC TAC AGT GCC TCT TTC CTC
      GTC GTT TTT GGT CCA TTT CGG GGT TTT GAG GAG TAG ATG TCA CGG AGA AAG GAG
      Q   Q   K   P   G   K   A   P   K   L   L   L   I   Y   S   A   S   F   L>

      120     130     140     150     160

170     TAT AGT GGT GTA CCA CCA GGT AAA GGT TTC AGC GGA TCC GGT AGT GGT ACT GAT TTC ACC
      ATA TCA CCA CAT GGT ATG TCC AAG TCG CCT AGG CCA TCA CCA TGA CTA AAG TGG
      Y   S   G   V   P   Y   R   F   S   G   S   G   S   G   T   D   F   T>

      180     190     200     210

220     CTC ACG ATC AGT AGC CTC CAG CCA GAA GAT TTC GCC ACT TAT TAC TGT CAA CAG
      GAG TGC TAG TCA TCG GAG GTC GGT CTT CTA AAG CGG TGA ATA ATG ACA GTT GTC
      L   T   I   S   S   L   Q   P   E   D   F   A   T   Y   Y   C   Q   Q>

      230     240     250     260     270

280     TAT AAC ATC TAC CCA CTC ACA TTC GGT CAG GGT ACT AAA GTA GAA ATC AAA
      ATA TTG TAG ATG GGT GAG TGT AAG CCA GTC CCA TGA TTT CAT CTT TAG TTT
      Y   N   I   Y   P   L   T   F   G   Q   G   T   K   V   E   I   K>

      290     300     310     320

```

8/27

FIG. 10 Grafted Vh sequence of hTNF40 (SEQ ID NO: 10)

```

10  CAG GTG CAG CTG GTC CAG TCA GGA GCA GAG GTT AAG AAG CCT GGT GCT TCC GTC
20  GTC CAC GTC GAC CAG GTC AGT CCT CGT CTC CAA TTC TTC GGA CCA CGA AGG CAG
30  Q V Q L V Q S G A E V K K P G A S V>
40
50
60  AAA GTT TCG TGT AAG GCC TCA GGC TAC GTG TTC ACA GAC TAT GGT ATG AAT TGG
70  TTT CAA AGC ACA TTC CGG AGT CCG ATG CAC AAG TGT CTG ATA CCA TAC TTA ACC
80  K V S C K A S G Y V F T D Y G M N W>
90
100
110 120 130 140 150 160
GTC AGA CAG GCC CCG GGA CAA GGC CTG GAA TGG ATG GGT TGG ATT AAT ACT TAC
CAG TCT GTC CGG GGC CCT GTT CCG GAC CTT ACC TAC CCA ACC TAA TTA TGA ATG
V R Q A P G Q G L E W M G W I N T Y>
170 180 190 200 210
ATT GGA GAG CCT ATT TAT TAT GCT CAA AAG TTC CAG GGC AGA GTC ACG TTC ACT CTA
TAA CCT CTC GGA TAA ATA CGA GTT TTC AAG GTC CCG TCT CAG TGC AAG TGA GAT
I G E P I Y A Q K F Q G R V T F T L>
220 230 240 250 260 270
GAC ACC TCC ACA AGC ACT GCA TAC ATG GAG CTG TCA TCT CTG AGA TCC GAG GAC
CTG TGG AGG TGT TCG TGA CGT ATG TAC CTC GAC AGT AGA GAC TCT AGG CTC CTG
D T S T S T A Y M E L S S L R S E D>
280 290 300 310 320
ACC GCA GTG TAC TAT TGT GCT AGA GGA TAC AGA TCT TAT GCC ATG GAC TAC TGG
TGG CGT CAC ATG ATA ACA CGA TCT CCT ATG TCT AGA ATA CGG TAC CTG ATG ACC
T A V Y Y C A R G Y R S Y A M D Y W>
330 340 350
GGC CAG GGT ACC CTA GTC ACA GTC TCC TCA
CCG GTC CCA TGG GAT CAG TGT CAG AGG AGT
G Q G T L V T V S S>

```

9/27

FIG. 11 Grafted Vh Sequence of hTNF40.4 (SEQ ID NO: 11)

```

10  GAG GTT CAG CTG GTC GAG TCA GGA GGC GGT CTC GTG CAG CCT GGC GGA TCA CTG
    CTC CAA GTC GAC CAG CTC AGT CCT CCG CCA GAG CAC GTC GGA CCG CCT AGT GAC
    E V Q L V E S G G L V Q P G G S L>
20
30
40
50
60  AGA TTG TCC TGT GCT GCA TCT GGT TAC GTC TTC ACA GAC TAT GGA ATG AAT TGG
    TCT AAC AGG ACA CGA CGT AGA CCA ATG CAG AAG TGT CTG ATA CCT TAC TTA ACC
    R L S C A A S G Y V F T D Y G M N W>
70
80
90
100
110 GTT AGA CAG GCC CCG GGA AAG GGC CTG GAA TGG ATG GGT TGG ATT AAT ACT TAC
    CAA TCT GTC CGG GGC CCT TTC CCG GAC CTT ACC TAC CCA ACC TAA TTA TGA ATG
    V R Q A P G K G L E W M G W I N T Y>
120
130
140
150
160
170 ATT GGA GAG CCT ATT TAT GCT GAC AGC GTC AAG GGC AGA TTC ACG TGC AAG AGA GAT
    TAA CCT CTC GGA TAA ATA CGA CTG TCG CAG TTC CCG TCT AAG TGC AAG AGA GAT
    I G E P I Y A D S V K G R F T F S L>
180
190
200
210
220 GAC ACA TCC AAG TCA ACA GCA TAC CTC CAA ATG AAT AGC CTG AGA GCA GAG GAC
    CTG TGT AGG TTC AGT TGT CGT ATG GAG GTT TAC TTA TCG GAC TCT CGT CTC CTG
    D T S K S T A Y L Q M N S L R A E D>
230
240
250
260
270
280 ACC GCA GTG TAC TAT TGT GCT AGA GGA TAC AGA TCT TAT GCC ATG GAC TAC TGG
    TGG CGT CAC ATG ATA ACA CGA TCT CCT ATG TCT AGA ATA CGG TAC TAC CTG ATG ACC
    T A V Y Y C A R G Y R S Y A M D Y W>
290
300
310
320
330 GGC CAG GGT ACC CTA GTC ACA GTC TCC TCA
    CCG GTC CCA TGG GAT CAG TGT CAG AGG AGT
    G Q G T L V T V S S>
340
350

```



10/27

FIG. 12







FIG. 13







12/27

## FIG. 15


Sequence of OmpA Oligonucleotide Adapter (SEQ ID NO: 101)



- Internal restriction sites are shown in bold
- The 5' XhoI cohesive end ligates into the Vector Sall site, blocking it
- S.D. represents the OmpA Shine Dalgarno sequence




14/27



15/27

FIG. 19



# FIG. 20 OLIGONUCLEOTIDE CASSETTES ENCODING DIFFERENT INTERGENIC SEQUENCES FOR E. Coli Fab' EXPRESSION

IGS CASSETTE-1; Intergenic space = -1

G, AGC, TCA, CCA, GTA, ACA, AAA, AGT, TTT, AAT, AGA, GGA, GAG, TGT, TAATG, AAG, AAG, ACT, GCT, ATA, GCA, ATT, G (SEQ ID No: 102)

S S P V T K S F N R G E C \* M K K T A I A I

End of c-Kappa sequence -> Start of OmpA sequence ->

IGS CASSETTE-2; Intergenic space = +1

G, AGC, TCA, CCA, GTA, ACA, AAA, AGT, TTT, AAT, AGA, GGG, GAG, TGT, TAA AATG, AAG, AAG, ACT, GCT, ATA, GCA, ATT, G (SEQ ID No: 103)

S S P V T K S F N R G E C \* M K K T A I A I

IGS CASSETTE-3; Intergenic space = +13

G, AGC, TCA, CCA, GTA, ACA, AAA, AGC, TTT, AAT, AGA, GGA, GAG, TGT, TGA GGAGGAAAAAAAATG, AAG, AAA, ACT, GCT, ATA, GCA, ATT, G (SEQ ID No: 104)


S S P V T K S F N R G E C \* M K K T A I A I

IGS CASSETTE-4; Intergenic space = +13

G, AGC, TCA, CCA, GTA, ACA, AAA, AGT, TTT, AAT, AGA, GGA, GAG, TGT, TGA CGAGGATTATATAATG, AAG, AAA, ACT, GCT, ATA, GCA, ATT, G (SEQ ID No: 105)


S S P V T K S F N R G E C \* M K K T A I A I

17/27






18/27




19/27

FIG. 23




20/27

FIG. 24




21/27

FIG. 24(contd.)



22/27

FIG. 24(contd.)



23/27

FIG. 24(contd.)




FIG. 24(contd.)






FIG. 24(contd.)






26/27

FIG. 24(contd.)



27/27

FIG. 24(contd.)



## SEQUENCE LISTING

<110> CELLTECH CHIROSCIENCE LIMITED

<120> BIOLOGICAL PRODUCTS

<130> P021741WO

<140>

<141>

<160> 115

<170> PatentIn Ver. 2.1

<210> 1

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: hTNF40 CDRH1

<400> 1

Asp Tyr Gly Met Asn  
1 5

<210> 2

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: hTNF40/human hybrid  
CDRH2

<400> 2

Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Ala Asp Ser Val Lys  
1 5 10 15

Gly

<210> 3

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: hTNF40 CDRH3

<400> 3

Gly Tyr Arg Ser Tyr Ala Met Asp Tyr  
1 5

<210> 4  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:hTNF40 CDRL1

<400> 4  
Lys Ala Ser Gln Asn Val Gly Thr Asn Val Ala  
1 5 10

<210> 5  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:hTNF40 CDRL2

<400> 5  
Ser Ala Ser Phe Leu Tyr Ser  
1 5

<210> 6  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:hTNF40 CDRL3

<400> 6  
Gln Gln Tyr Asn Ile Tyr Pro Leu Thr  
1 5

<210> 7  
<211> 17  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:hTNF40 CDRH2

<400> 7  
Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Val Asp Asp Phe Lys  
1 5 10 15

Gly

<210> 8

<211> 321  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)..(321)

<220>  
 <223> Description of Artificial Sequence:hTNF40-gL1

<400> 8  
 gac att caa atg acc cag agc cca tcc agc ctg agc gca tct gta gga 48  
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
 1 5 10 15  
 gac cgg gtc acc atc act tgt aaa gcc agt cag aac gta ggt act aac 96  
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asn Val Gly Thr Asn  
 20 25 30  
 gta gcc tgg tat cag caa aaa cca ggt aaa gcc cca aaa gcc ctc atc 144  
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Ala Leu Ile  
 35 40 45  
 tac agt gcc tct ttc ctc tat agt ggt gta cca tac agg ttc agc gga 192  
 Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Tyr Arg Phe Ser Gly  
 50 55 60  
 tcc ggt agt ggt act gat ttc acc ctc acg atc agt agc ctc cag cca 240  
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
 65 70 75 80  
 gaa gat ttc gcc act tat tac tgt caa cag tat aac atc tac cca ctc 288  
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ile Tyr Pro Leu  
 85 90 95  
 aca ttc ggt cag ggt act aaa gta gaa atc aaa 321  
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
 100 105

<210> 9  
 <211> 321  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)..(321)

<220>  
 <223> Description of Artificial Sequence:hTNF40-gL2

<400> 9  
 gac att caa atg acc cag agc cca tcc agc ctg agc gca tct gta gga 48  
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

```

1           5           10           15
gac cgg gtc acc atc act tgt aaa gcc agt cag aac gta ggt act aac 96
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asn Val Gly Thr Asn
                20                25                30

gta gcc tgg tat cag caa aaa cca ggt aaa gcc cca aaa ctc ctc atc 144
Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
                35                40                45

tac agt gcc tct ttc ctc tat agt ggt gta cca tac agg ttc agc gga 192
Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Tyr Arg Phe Ser Gly
                50                55                60

tcc ggt agt ggt act gat ttc acc ctc acg atc agt agc ctc cag cca 240
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
                65                70                75                80

gaa gat ttc gcc act tat tac tgt caa cag tat aac atc tac cca ctc 288
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ile Tyr Pro Leu
                85                90                95

aca ttc ggt cag ggt act aaa gta gaa atc aaa 321
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
                100                105

<210> 10
<211> 354
<212> DNA
<213> Artificial Sequence

<220>
<221> CDS
<222> (1)..(354)

<220>
<223> Description of Artificial Sequence:ghlhtNF40.4 (Figure 10)

<400> 10
cag gtg cag ctg gtc cag tca gga gca gag gtt aag aag cct ggt gct 48
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1           5           10           15

tcc gtc aaa gtt tcg tgt aag gcc tca ggc tac gtg ttc aca gac tat 96
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Val Phe Thr Asp Tyr
                20                25                30

ggt atg aat tgg gtc aga cag gcc ccg gga caa ggc ctg gaa tgg atg 144
Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
                35                40                45

ggt tgg att aat act tac att gga gag cct att tat gct caa aag ttc 192
Gly Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Ala Gln Lys Phe
                50                55                60

```

```

cag ggc aga gtc acg ttc act cta gac acc tcc aca agc act gca tac 240
Gln Gly Arg Val Thr Phe Thr Leu Asp Thr Ser Thr Ser Thr Ala Tyr
 65                70                75                80

atg gag ctg tca tct ctg aga tcc gag gac acc gca gtg tac tat tgt 288
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
      85                90                95

gct aga gga tac aga tct tat gcc atg gac tac tgg ggc cag ggt acc 336
Ala Arg Gly Tyr Arg Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr
      100                105                110

cta gtc aca gtc tcc tca 354
Leu Val Thr Val Ser Ser
      115

```

<210> 11  
 <211> 354  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)..(354)

<220>  
 <223> Description of Artificial Sequence:gh3hTNF40.4 (Figure 11)

```

<400> 11
gag gtt cag ctg gtc gag tca gga ggc ggt ctc gtg cag cct ggc gga 48
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1                5                10                15

tca ctg aga ttg tcc tgt gct gca tct ggt tac gtc ttc aca gac tat 96
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Val Phe Thr Asp Tyr
      20                25                30

gga atg aat tgg gtt aga cag gcc ccg gga aag ggc ctg gaa tgg atg 144
Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met
      35                40                45

ggt tgg att aat act tac att gga gag cct att tat gct gac agc gtc 192
Gly Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Ala Asp Ser Val
      50                55                60

aag ggc aga ttc acg ttc tct cta gac aca tcc aag tca aca gca tac 240
Lys Gly Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
      65                70                75                80

ctc caa atg aat agc ctg aga gca gag gac acc gca gtg tac tat tgt 288
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
      85                90                95

gct aga gga tac aga tct tat gcc atg gac tac tgg ggc cag ggt acc 336

```

Ala Arg Gly Tyr Arg Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr  
 100 105 110

cta gtc aca gtc tcc tca  
 Leu Val Thr Val Ser Ser  
 115

354

<210> 12  
 <211> 9  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:part of a primer sequence

<400> 12  
 gccgccacc

9

<210> 13  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer CH1

<400> 13  
 atgaaatgca gctgggtcat sttctt

26

<210> 14  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer CH2

<400> 14  
 atgggatgga gctratatcat sytctt

26

<210> 15  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer CH3

<400> 15  
 atgaagwtgt gggttaaactg ggtttt

26



<210> 16  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH4

<400> 16  
atgractttg ggytcagctt grt 23

<210> 17  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH5

<400> 17  
atggactcca ggctcaattt agtttt 26

<210> 18  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH6

<400> 18  
atggctgtcy trgsgctrct cttctg 26

<210> 19  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH7

<400> 19  
atggratgga gckggrtctt tmtctt 25

<210> 20  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH8

<400> 20

atgagagtgc tgattctttt gtg

23

<210> 21  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH9

<400> 21  
atggmttggg tgtggamctt gctatt

26

<210> 22  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH10

<400> 22  
atgggcagac ttacattctc attcct

26

<210> 23  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH11

<400> 23  
atggattttg ggctgatttt ttttattg

28

<210> 24  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CH12

<400> 24  
atgatggtgt taagtctttct gtacct

26

<210> 25  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer 5' end

<400> 25

gcgcgcaagc ttgccgccac c 21

<210> 26  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL1

<400> 26  
atgaagttgc ctgtaggct gttggtgct 29

<210> 27  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL2

<400> 27  
atggagwcag acacactcct gytatgggt 29

<210> 28  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL3

<400> 28  
atgagtgtgc tcactcaggt cct 23

<210> 29  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL4

<400> 29  
atgaggrccc ctgctcagwt tyttgg 26

<210> 30  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL5

<400> 30  
atggatttwc aggtgcagat twtcagctt 29

<210> 31  
<211> 29  
<212> DNA  
<213> Artificial Sequence

---

<220>  
<223> Description of Artificial Sequence:primer CL5A

<400> 31  
atggatttc argtcagat twtcagctt

29

<210> 32  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL6

<400> 32  
atgaggtkcy ytgytsagyt yctgrg

26

<210> 33  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL7

<400> 33  
atgggcwtca agatggagtc aca

23

<210> 34  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL8

<400> 34  
atgtggggay ctktttycmm tttttcaat

29

<210> 35  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL9

<400> 35  
atggtrtccw casctcagtt cctt

24

<210> 36  
<211> 26

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL10

<400> 36  
atgtatatat gtttggtgtc tatttc 26

<210> 37  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL11

<400> 37  
atggaagccc cagctcagct tctctt 26

<210> 38  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL12A

<400> 38  
atgragtywc agaccaggt cttyrt 26

<210> 39  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL12B

<400> 39  
atggagacac attctcaggt ctttgt 26

<210> 40  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer CL13

<400> 40  
atggattcac aggcccaggt tcttat 26

<210> 41  
<211> 26  
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL14

<400> 41

atgatgagtc ctgcccagtt cctggt

26

<210> 42

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL15

<400> 42

atgaatttgc ctgttcacat cttggtgct

29

<210> 43

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL16

<400> 43

atggattttc aattggtcct catctcctt

29

<210> 44

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL17A

<400> 44

atgaggtgcc tarcttsagtt cctgrg

26

<210> 45

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL17B

<400> 45

atgaagtact ctgctcagtt tctagg

26

<210> 46

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL17C

<400> 46

atgaggcatt ctcttcaatt cttggg

26

<210> 47

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer 5' end

<400> 47

ggactgttcg aagccgccac c

21

<210> 48

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer CL12

<400> 48

ggatacagtt ggtgcagcat ccgtacgttt

30

<210> 49

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer R2155

<400> 49

gcagatgggc ccttcggtga ggctgmrgag acdgtga

37

<210> 50

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer R1053

<400> 50

gctgacagac taacagactg ttcc

24

<210> 51

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer R720

<400> 51  
gctctcggag gtgctcct

18

<210> 52  
<211> 70  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:oligonucleotide P7982

<400> 52  
gaattcaggg tcaccatcac ttgtaaagcc agtcagaacg taggtactaa cgtagcctgg 60  
tatacagcaaa 70

<210> 53  
<211> 71  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:oligonucleotide P7983

<400> 53  
atagaggaaa gaggcactgt agatgagggc ttttggggct ttacctgggt tttgctgata 60  
ccaggctacg t 71

<210> 54  
<211> 71  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:oligonucleotide P7984

<400> 54  
tacagtgcct ctttcctcta tagtggtgta ccatacaggt tcagcggatc cggtagtgggt 60  
actgatttca c 71

<210> 55  
<211> 71  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:oligonucleotide P7985

<400> 55  
gacagtaata agtggcgaaa tcttctgggt ggaggctact gatcgtgagg gtgaaatcag 60  
taccactacc g 71

<210> 56  
<211> 89  
<212> DNA  
<213> Artificial Sequence



&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide P7986

&lt;400&gt; 56

atttcgccac ttattactgt caacagtata acatctaccc actcacattc ggtcagggta 60  
ctaaagtaga aatcaaacgt acggaattc 89

&lt;210&gt; 57

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide P7981

&lt;400&gt; 57

gaattcaggg tcaccatcac ttgtaaagcc 30

&lt;210&gt; 58

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide P7980

&lt;400&gt; 58

gaattccgta cgtttgattt ctactttagt 30

&lt;210&gt; 59

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide R1053

&lt;400&gt; 59

gctgacagac taacagactg ttcc 24

&lt;210&gt; 60

&lt;211&gt; 57

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide R5350

&lt;400&gt; 60

tctagatggc acaccatctg ctaagtttga tgcagcatag atcaggagct taggagc 57

&lt;210&gt; 61

&lt;211&gt; 59

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide R5349

&lt;400&gt; 61

gcagatggtg tgccatctag attcagtggc agtggatcag gcacagactt taccctaac 59

&lt;210&gt; 62

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:oligonucleotide R684

&lt;400&gt; 62

ttcaactgct catcagat

18

&lt;210&gt; 63

&lt;211&gt; 65

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer P7989

&lt;400&gt; 63

gaagcaccag gcttcttaac ctctgctcct gactggacca gctgcacctg agagtgcacg 60  
aattc 65

&lt;210&gt; 64

&lt;211&gt; 71

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer P7990

&lt;400&gt; 64

ggttaagaag cctggtgctt ccgtcaaagt ttcgtgtaag gcctcaggct acgtgttcac 60  
agactatggt a 71

&lt;210&gt; 65

&lt;211&gt; 71

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer P7991

&lt;400&gt; 65

ccaacccatc catttcaggc cttgtccogg ggcctgcttg acccaattca taccatagtc 60  
tgtgaacacg t 71

&lt;210&gt; 66

&lt;211&gt; 81

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7995

<400> 66

ggcctgaaat ggatgggttg gattaatact tacattggag agcctattta tgttgacgac 60  
ttcaagggca gattcacgtt c 81

<210> 67

<211> 56

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7992

<400> 67

ccatgtatgc agtgcgttgt ggaggtgtct agagtgaacg tgaatctgcc cttgaa 56

<210> 68

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7993

<400> 68

ccacaagcac tgcatacatg gagctgtcat ctctgagatc cgaggacacc gcagtgtact 60  
at 62

<210> 69

<211> 78

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7994

<400> 69

gaattcggta ccctggcccc agtagtccat ggcataagat ctgtatcctc tagcacaata 60  
gtacactgcg gtgtcctc 78

<210> 70

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7988

<400> 70

gaattcgtgc actctcaggt gcagctggtc 30

<210> 71

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7987

<400> 71

gaattcggta ccctggcccc agtagtccat

30

<210> 72

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7999

<400> 72

gatccgccag gctgcacgag accgcctcct gactcgacca gctgaacctc agagtgcacg 60  
aatc 65

<210> 73

<211> 71

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P8000

<400> 73

tctcgtgcag cctggcggat cgctgagatt gtccgtgtgt gcatctggtt acgtcttcac 60  
agactatgga a 71

<210> 74

<211> 71

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P8001

<400> 74

ccaacccatc catttcaggc cctttcccgg ggcctgctta acccaattca ttccatagtc 60  
tgtgaagacg t 71

<210> 75

<211> 55

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7997

<400> 75

ggaggtatgc tggtgacttg gatgtgtcta gagagaacgt gaatctgccc ttgaa 55

<210> 76

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7998

<400> 76

ccaagtcaac agcatacctc caaatgaata gcctgagagc agaggacacc gcagtgtact 60  
at 62

<210> 77

<211> 78

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7993

<400> 77

gaattcggtta ccctggcccc agtagtccat ggcataagat ctgtatcctc tagcacaata 60  
gtacactgcg gtgtcctc 78

<210> 78

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer P7996

<400> 78

gaattcgtgc actctgaggt tcagctggtc 30

<210> 79

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 5' primer

<400> 79

cgcgcggcaa ttgcagtggc cttggctggt ttogctaccg tagcgcaagc tgacattcaa 60  
atgacccaga gcc 74

<210> 80

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 3' primer

<400> 80

ttcaactgct catcagatgg 20

<210> 81  
 <211> 78  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: 5' primer

<400> 81  
 gctatcgcaa ttgcagtggc gctagctggt ttgccaccg tggcgcaagc tgaggttcag 60  
 ctggtcgagt caggaggc 78

<210> 82  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: 3' primer

<400> 82  
 gcctgagttc cacgacac

18

<210> 83  
 <211> 23  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: human group 1 consensus  
 framework L1

<400> 83  
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly  
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys  
 20

<210> 84  
 <211> 23  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: hTNF40 framework L1

<400> 84  
 Asp Ile Val Met Thr Gln Ser Gln Lys Phe Met Ser Thr Ser Val Gly  
 1 5 10 15

Asp Arg Val Ser Val Thr Cys  
 20

<210> 85  
 <211> 15  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:human group 1 consensus  
 framework L2

<400> 85  
 Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr  
     1                    5                    10                    15

<210> 86  
 <211> 15  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:hTNF40 framework L2

<400> 86  
 Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Ala Leu Ile Tyr  
     1                    5                    10                    15

<210> 87  
 <211> 32  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:human group 1 consensus  
 framework L3

<400> 87  
 Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr  
     1                    5                    10                    15

Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys  
                     20                    25                    30

<210> 88  
 <211> 32  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:hTNF40 framework L3

<400> 88  
 Gly Val Pro Tyr Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr  
     1                    5                    10                    15

Leu Thr Ile Ser Thr Val Gln Ser Glu Asp Leu Ala Glu Tyr Phe Cys  
                   20                                  25                                  30

<210> 89

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:human group 1 consensus  
                   framework L4

<400> 89

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg  
   1                                  5                                  10

<210> 90

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:hTNF40 framework L4

<400> 90

Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg  
   1                                  5                                  10

<210> 91

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:human group 1 consensus  
                   framework H1

<400> 91

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
   1                                  5                                  10                                  15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr  
                   20                                  25                                  30

<210> 92

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:hTNF40 framework H1



&lt;400&gt; 92

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu  
 1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Val Phe Thr  
 20 25 30

&lt;210&gt; 93

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:human group 1 consensus framework H2

&lt;400&gt; 93

Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met Gly  
 1 5 10

&lt;210&gt; 94

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:hTNF40 framework H2

&lt;400&gt; 94

Trp Val Lys Gln Ala Pro Gly Lys Ala Phe Lys Trp Met Gly  
 1 5 10

&lt;210&gt; 95

&lt;211&gt; 32

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:human group 1 consensus framework H3

&lt;400&gt; 95

Arg Val Thr Ile Thr Arg Asp Thr Ser Thr Ser Thr Ala Tyr Met Glu  
 1 5 10 15

Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg  
 20 25 30

&lt;210&gt; 96

<211> 32  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:hTNF40 framework H3

<400> 96  
 Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Phe Leu Gln  
           1                          5                          10                          15  
 Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys Ala Arg  
                           20                          25                          30

<210> 97  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:human group 1 consensus  
           framework H4

<400> 97  
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
           1                          5                          10

<210> 98  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:hTNF40 framework H4

<400> 98  
 Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
           1                          5                          10

<210> 99  
 <211> 324  
 <212> DNA  
 <213> murine

<220>  
 <221> CDS  
 <222> (1)..(324)  
 <223> mouse hTNF40 light chain variable domain

<400> 99

```

gac att gtg atg acc cag tct caa aaa ttc atg tcc aca tca gta gga 48
Asp Ile Val Met Thr Gln Ser Gln Lys Phe Met Ser Thr Ser Val Gly
  1             5             10             15

gac agg gtc agc gtc acc tgc aag gcc agt cag aat gtg ggt act aat 96
Asp Arg Val Ser Val Thr Cys Lys Ala Ser Gln Asn Val Gly Thr Asn
          20             25             30

gta gcc tgg tat caa cag aaa cca gga caa tct cct aaa gca ctg att 144
Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Ala Leu Ile
          35             40             45

tac tgc gca tcc ttc cta tat agt gga gtc cct tat cgc ttc aca ggc 192
Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Tyr Arg Phe Thr Gly
          50             55             60

agt gga tct ggg aca gat ttc act ctc acc atc agc act gtg cag tct 240
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Thr Val Gln Ser
  65             70             75             80

gaa gac ttg gca gag tat ttc tgt cag caa tat aac atc tat cct ctc 288
Glu Asp Leu Ala Glu Tyr Phe Cys Gln Gln Tyr Asn Ile Tyr Pro Leu
          85             90             95

acg ttc ggt gct ggg acc aag ctg gag ctg aaa cgt 324
Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg
          100             105

```

&lt;210&gt; 100

&lt;211&gt; 354

&lt;212&gt; DNA

&lt;213&gt; murine

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(354)

&lt;223&gt; mouse hTNF40 heavy chain variable domain

&lt;400&gt; 100

```

cag atc cag ttg gtg cag tct gga cct gag ctg aag aag cct gga gag 48
Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
  1             5             10             15

aca gtc aag atc tcc tgc aag gct tct gga tat gtt ttc aca gac tat 96
Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Val Phe Thr Asp Tyr
          20             25             30

gga atg aat tgg gtg aag cag gct cca gga aag gct ttc aag tgg atg 144
Gly Met Asn Trp Val Lys Gln Ala Pro Gly Lys Ala Phe Lys Trp Met
          35             40             45

ggc tgg ata aac acc tac att gga gag cca ata tat gtt gat gac ttc 192
Gly Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Val Asp Asp Phe
          50             55             60

```

aag gga cga ttt gcc ttc tct ttg gaa acc tct gcc agc act gcc ttt 240  
 Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Phe  
 65 70 75 80

ttg cag atc aac aac ctc aaa aat gag gac acg gct aca tat ttc tgt 288  
 Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys  
 85 90 95

gca aga ggt tac cgg tcc tat gct atg gac tac tgg ggt caa gga acc 336  
 Ala Arg Gly Tyr Arg Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr  
 100 105 110

tca gtc acc gtc tct tca  
 Ser Val Thr Val Ser Ser 354  
 115

<210> 101

<211> 84

<212> DNA

<213> Artificial Sequence

<220>

<221> CDS

<222> (29)..(67)

<223> Description of Artificial Sequence:OmpA oligonucleotide adaptor

<400> 101

tcgagttcta gataacgagg cgtaaaaa atg aaa aag aca gct atc gca att 52  
 Met Lys Lys Thr Ala Ile Ala Ile  
 1 5

gca gtg gcc ttg gct ctgacgtacg agtcagg 84  
 Ala Val Ala Leu Ala  
 10

<210> 102

<211> 67

<212> DNA

<213> Artificial Sequence

<220>

<221> CDS

<222> (2)..(40)

<220>

<221> CDS

<222> (43)..(66)

<220>

<223> Description of Artificial Sequence:IGS cassette-1

<400> 102

g agc tca cca gta aca aaa agt ttt aat aga gga gag tgt ta atg aag 48

Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Xaa	Xaa	Lys
1				5					10					15

aag act gct ata gca att g 67  
 Lys Thr Ala Ile Ala Ile  
 20

<210> 103  
 <211> 69  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (2)..(43)

<220>  
 <221> CDS  
 <222> (45)..(68)

<220>  
 <223> Description of Artificial Sequence:IGS cassette-2

<400> 103	
g agc tca cca gta aca aaa agt ttt aat aga ggg gag tgt taa a atg	47
Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys Met	
1 5 10 15	

aag aag act gct ata gca att g 69  
 Lys Lys Thr Ala Ile Ala Ile  
 20

<210> 104  
 <211> 81  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (2)..(43)

<220>  
 <221> CDS  
 <222> (57)..(80)

<220>  
 <223> Description of Artificial Sequence:IGS cassette-3

<400> 104	
g agc tca cca gta aca aaa agc ttt aat aga gga gag tgt tga	43
Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys	
1 5 10	

ggaggaaaaa aaa atg aag aaa act gct ata gca att g 81  
 Met Lys Lys Thr Ala Ile Ala Ile

15

20

<210> 105  
 <211> 81  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (2)..(43)

<220>  
 <221> CDS  
 <222> (57)..(80)

<220>  
 <223> Description of Artificial Sequence:IGS cassette-4

<400> 105  
 g agc tca cca gta aca aaa agt ttt aat aga gga gag tgt tga 43  
 Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
 1 5 10

cgaggattat ata atg aag aaa act gct ata gca att g 81  
 Met Lys Lys Thr Ala Ile Ala Ile  
 15 20

<210> 106  
 <211> 30  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:human group 3 consensus  
 framework H1

<400> 106  
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser  
 20 25 30

<210> 107  
 <211> 13  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:human group 3 consensus  
 framework H2

<400> 107

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ser  
 1 5 10

<210> 108

<211> 32

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:human group 3 consensus  
 framework H3

<400> 108

Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu Gln  
 1 5 10 15

Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg  
 20 25 30

<210> 109

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:human group 3 consensus  
 framework H4

<400> 109

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
 1 5 10

<210> 110

<211> 648

<212> DNA

<213> Artificial Sequence

<220>

<223> Grafted Heavy Chain for Fab

<400> 110

gaggttcagc	tggtcgagtc	aggaggcgg	ctcgtgcagc	ctggcggatc	actgagattg	60
tcctgtgctg	catctgggta	cgtcttcaca	gactatggaa	tgaattgggt	tagacaggcc	120
ccgggaaagg	gcctgggaat	gatgggttgg	attaatactt	acattggaga	gcctatztat	180
gctgacagcg	tcaagggcag	attcacgttc	tctctagaca	catccaagtc	aacagcatac	240
ctccaaatga	atagcctgag	agcagaggac	accgcagtgt	actattgtgc	tagaggatac	300
agatcttatg	ccatggacta	ctggggccag	ggtaccctag	tcacagtctc	ctcagcttcc	360
accaagggcc	catcgggtct	ccccctggca	ccctcctcca	agagcacctc	tgggggcaca	420
gcggccctgg	gctgcctgg	caaggactac	ttccccgaac	cggtgacgg	gtcgtggaac	480
tcaggcgccc	tgaccagcgg	cgtgcacacc	ttcccggtg	tcctacagtc	ctcaggactc	540
tactccctca	gcagcgtgg	gaccgtgccc	tccagcagct	tgggcaccca	gacctacatc	600

tgcaacgtga atcacaagcc cagcaacacc aaggtcgaca agaaagtt

648

<210> 111

<211> 216

<212> PRT

<213> Artificial Sequence

<220>

<223> Grafted Heavy Chain for Fab

<400> 111

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Val Phe Thr Asp Tyr  
20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met  
35 40 45

Gly Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Ala Asp Ser Val  
50 55 60

Lys Gly Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr  
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Gly Tyr Arg Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro  
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly  
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn  
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln  
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser  
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser  
195 200 205

Asn Thr Lys Val Asp Lys Lys Val  
210 215

<210> 112

<211> 642

<212> DNA

<213> Artificial Sequence



&lt;220&gt;

&lt;223&gt; Grafted Light Chain for Fab and Modified Fab

&lt;400&gt; 112

```

gacattcaaa tgaccagag cccatccagc ctgagcgcat ctgtaggaga ccgggtcacc 60
atcacttgta aagccagtca gaacgtaggt actaacgtag cctggatatca gcaaaaacca 120
ggtaaagccc caaaagccct catctacagt gcctctttcc tctatagtgg tgtaccatac 180
aggttcagcg gatccggtag tggtagctgat ttcaccctca cgatcagtag cctccagcca 240
gaagatttcg ccacttatta ctgtcaacag tataacatct acccactcac attcggtcag 300
ggtactaaag tagaaatcaa acgtacggta ggggccccat ctgtcttcat cttcccgcca 360
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taacttctat 420
cccagagagg ccaaagtaca gtggaagggtg gataacgccc tccaatoggg taactcccag 480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg 540
ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc 600
ctgagctcac cagtaacaaa aagctttaat agaggagagt gt 642

```

&lt;210&gt; 113

&lt;211&gt; 214

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Grafted Light Chain for Fab and Modified Fab

&lt;400&gt; 113

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asn Val Gly Thr Asn
20           25           30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Ala Leu Ile
35           40           45

Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Tyr Arg Phe Ser Gly
50           55           60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ile Tyr Pro Leu
85           90           95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
100          105          110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115          120          125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130          135          140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145          150          155          160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser

```

165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr  
180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser  
195 200 205

Phe Asn Arg Gly Glu Cys  
210

<210> 114  
<211> 687  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Grafted Heavy Chain for Modified Fab

<400> 114

gaggttcagc	tggtcgagtc	aggaggcggt	ctcgtgcagc	ctggcggatc	actgagattg	60
tcctgtgctg	catctgggta	cgtcttcaca	gactatggaa	tgaattgggt	tagacaggcc	120
ccgggaaagg	gcctggaatg	gatgggttgg	attaatactt	acattggaga	gcctatttat	180
gctgacagcg	tcaagggcag	attcacgttc	tctctagaca	catccaagtc	aacagcatac	240
ctccaaatga	atagcctgag	agcagaggac	accgcagtgt	actattgtgc	tagaggatac	300
agatcttatg	ccatggacta	ctggggccag	ggtaccctag	tcacagtctc	ctcagcttcc	360
accaagggcc	catcggtctt	ccccctggca	ccctcctcca	agagcacctc	tgggggcaca	420
gcggccctgg	gctgcctggg	caaggactac	ttccccgaac	cggtgacggt	gtcgtggaac	480
tcaggcgccc	tgaccagcgg	cgtgcacacc	ttcccgctg	tcctacagtc	ctcaggactc	540
tactccctca	gcagcgtggg	gaccgtgccc	tccagcagct	tgggcaccca	gacctacatc	600
tgcaacgtga	atcacaagcc	cagcaacacc	aaggtcgaca	agaaagtga	gcccaaactc	660
tgtgacaaaa	ctcacacatg	cgccgcg				687

<210> 115  
<211> 229  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Grafted Heavy Chain for Modified Fab

<400> 115

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	1	5	10	15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Val Phe Thr Asp Tyr	20	25	30	
Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met	35	40	45	
Gly Trp Ile Asn Thr Tyr Ile Gly Glu Pro Ile Tyr Ala Asp Ser Val	50	55	60	
Lys Gly Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr	65	70	75	80

Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys		
				85					90					95			
Ala	Arg	Gly	Tyr	Arg	Ser	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr		
				100					105					110			
Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro		
				115					120					125			
Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly		
				130					135					140			
Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn		
145					150					155					160		
Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln		
				165					170					175			
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser		
				180					185					190			
Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser		
				195					200					205			
Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr		
				210					215					220			
His	Thr	Cys	Ala	Ala													
225																	

# INTERNATIONAL SEARCH REPORT

In International Application No  
PCT/GB 01/02477

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/13 C07K16/24 C07K16/46 A61K47/48 C07K19/00  
C12N15/62 C12N15/70 C12N1/21 A61K39/395 A61P19/02  
A61P37/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched - (classification system followed by classification symbols)  
IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)  
BIOSIS, EMBASE, CHEM ABS Data, WPI Data, PAJ, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 25971 A (CELLTECH THERAPEUTICS LIMITED) 18 June 1998 (1998-06-18)	1-4, 13, 15, 16, 26, 28-32, 60-64, 67
Y	examples 2, 5, 6 claims	6-12, 14, 17-23, 27, 33-41, 49-54, 56-59
	---	
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

15 August 2001

Date of mailing of the international search report

22/08/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Nooij, F

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/02477

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 92 11383 A (CELLTECH LIMITED) 9 July 1992 (1992-07-09)  examples claims  ---	6-12, 14, 17-23, 27, 33-41, 49-54, 56-59
X	WO 99 64460 A (CELLTECH THERAPEUTICS LIMITED) 16 December 1990 (1990-12-16)  examples 3,4 claims  ---	1-4, 13, 15, 16, 26, 28-32, 60-64, 67
X	EP 0 380 068 A (MOLECULAR THERAPEUTICS, INC.) 1 August 1990 (1990-08-01) figure 4  ---	67
A	S. STEPHENS ET AL.: "Comprehensive pharmacokinetics of a humanized antibody and analysis of residual anti-idiotypic responses." IMMUNOLOGY, vol. 85, no. 4, August 1995 (1995-08), pages 668-674, XP000881488 Oxford, GB abstract page 668, right-hand column figure 1  ---	1-67
A	K. NAGAHIRA ET AL.: "Humanization of a mouse neutralizing monoclonal antibody against tumor necrosis factor-alpha (TNF-alpha)." JOURNAL OF IMMUNOLOGICAL METHODS, vol. 222, no. 1-2, 1 January 1999 (1999-01-01), pages 83-92, XP004152430 Amsterdam, The Netherlands abstract figure 2  -----	1-67

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/02477

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9825971	A	18-06-1998	AU 733251 B	10-05-2001
			AU 5404998 A	03-07-1998
			EP 0948544 A	13-10-1999
			GB 2334957 A, B	08-09-1999
WO 9211383	A	09-07-1992	AT 137534 T	15-05-1996
			AT 134387 T	15-03-1996
			AU 669083 B	23-05-1996
			AU 7772394 A	09-03-1995
			AU 651984 B	11-08-1994
			AU 8200591 A	04-02-1992
			AU 657937 B	30-03-1995
			AU 9108491 A	22-07-1992
			BG 60462 B	28-04-1995
			BR 9007197 A	28-01-1992
			BR 9106232 A	30-03-1993
			CA 2065325 A	06-01-1992
			CA 2076540 A	22-06-1992
			CA 2129554 A	22-06-1992
			DE 4193302 C	24-08-2000
			DE 4193302 T	18-02-1993
			DE 69022982 D	16-11-1995
			DE 69022982 T	28-03-1996
			DE 69117284 D	28-03-1996
			DE 69117284 T	05-09-1996
			DE 69119211 D	05-06-1996
			DE 69119211 T	19-12-1996
			DK 516785 T	18-03-1996
			EP 0460167 A	11-12-1991
			EP 0491031 A	24-06-1992
			EP 0516785 A	09-12-1992
			EP 0626389 A	30-11-1994
			EP 0927758 A	07-07-1999
			ES 2084338 T	01-05-1996
			FI 923737 A	20-08-1992
			GB 2246570 A, B	05-02-1992
			GB 2251859 A, B	22-07-1992
			GB 2257145 A, B	06-01-1993
			GB 2276169 A	21-09-1994
			GB 2279077 A, B	21-12-1994
			GR 3017734 T	31-01-1996
			GR 3019066 T	31-05-1996
			HU 62661 A	28-05-1993
			HU 215383 B	28-03-2000
			HU 9500283 A	28-12-1995
			JP 10136986 A	26-05-1998
			JP 3145401 B	12-03-2001
			JP 5502587 T	13-05-1993
			KR 253426 B	15-04-2000
			NL 9120013 T	02-11-1992
			NO 913228 A	21-10-1991
			NO 923231 A	20-10-1992
WO 9964460	A	16-12-1999	AU 4278399 A	30-12-1999
			DE 19983347 T	28-06-2001
			EP 1090037 A	11-04-2001
			GB 2354242 A	21-03-2001

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/02477

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 380068 A	01-08-1990	AT 81153 T	15-10-1992
		AU 641907 B	07-10-1993
		AU 4876690 A	02-08-1990
		CA 2008259 A	24-07-1990
		DE 69000338 D	05-11-1992
		DE 69000338 T	25-02-1993
		DK 380068 T	02-11-1992
		ES 2052077 T	01-07-1994
		JP 2295487 A	06-12-1990
		NO 900092 A	25-07-1990
		NZ 232201 A	28-04-1992
		PT 92900 A	31-07-1990
		ZA 9000468 A	31-10-1990

---

